

A

EXPRESS MAIL CERTIFICATE

Date 6/8/99 Label No. 9595859379  
I hereby certify that, on the date indicated above, I  
deposited this paper or fee with the U.S. Postal Service  
and that it was addressed for delivery to the Commissioner  
of Patents & Trademarks, Washington, DC 20231 by Express  
Mail Post Office to "Addressee" service.

Name (Print)

Signature

D B Beck [Signature]

JCS18 U.S. PTO  
09/32/563  
06/08/99

Docket No. 0776/1F216-US2

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re Application of:

Allan **SVENDSEN**, *et al.*

For: **α-AMYLASE MUTANTS**

CONTINUING APPLICATION

Hon. Commissioner of  
Patents and Trademarks  
Washington, DC 20231

Sir:

This is a request for filing under 37 C.F.R. 1.53(b) of a:

☒ Continuation ☐ Divisional

application of the following pending prior application:

Serial No. 08/683,838 Filed: July 18, 1996

Of: Allan **SVENDSEN**, *et al.*

For: **α-AMYLASE MUTANTS**

Examiner: E. Slobodyansky Group: 1652

1. Enclosed is a copy of the prior application as originally filed (along with a copy of the original Declaration). No amendments identified in the declaration for this prior application introduced new matter. Please use this copy as the application and declaration for the present case.

2. The filing fee is calculated below:

CLAIMS AS FILED, AFTER ACCOMPANYING AMENDMENT

	Claims on File	Number Extra	Rate
Basic Fee . . . . .			\$760.00
Total Claims . . . . .	43 - 20 =	23 x \$18	\$414.00
Independent Claims . . . . .	28 - 3 =	25 x \$78	\$1,950.00
If Multiple Dependent Claims Are Present, Add \$260.00 . . . . .			\$
Total Filing Fee . . . . .			\$3,124.00
For Small Entity (half of preceding total)* . . . . .			\$

\*No. 9 below must be checked to claim this reduction.

3. A check in the amount of \$3,124.00 is enclosed.
4. ☒ Cancel claims 1-69.
5. ☒ Amend the specification by inserting before the first line the sentence (check one and fill in):

"This is a ☒ continuation, ☐ division, of application Serial No. 08/683,838, filed July 18, 1996, which is a continuation-in-part of 08/600,908, filed February 13, 1996, which is a 371 of PCT/DK96/00057, filed February 5, 1996. Each of these prior applications is hereby incorporated herein by reference, in its entirety."

6. [X] The prior application is assigned to: Novo Nordisk A/S
7. [X] A Preliminary Amendment is also enclosed.
8. [X] Informal Drawings are filed herewith
9. [X] Associate Power of Attorney
10. [] A verified statement claiming small entity status (check one):  
[] was filed in parent application or [] is enclosed.
11. [X] Priority is claimed from  
Country: Denmark  
Number: 0128/95  
Date: February 3, 1995;  
  
Country: Denmark  
Number: 1192/95  
Date: October 23, 1995; and  
  
Country: Denmark  
Number: 1256/95  
Date: November 10, 1995

The priority documents

- [X] were filed in the prior applications. Certified copies of the priority applications were submitted in PCT/DK96/00057.
- [] is enclosed.

12. The Assignment

- ☒ was recorded at Reel 8217 Frame 0630 on November 12, 1996.  
☒ copy is enclosed.

Respectfully submitted,



Adda C. Gogoris

Reg. No. 29,714

Attorney for Applicant per Attached  
Associate Power of Attorney from  
Attorney

DARBY & DARBY P.C.  
805 Third Avenue  
New York, NY 10022  
212-527-7700

09327563-060699

EXPRESS MAIL CERTIFICATE

Date 6/8/99 Label No. cm595859379

I hereby certify that, on the date indicated above I deposited this paper or fee with the U.S. Postal Service and that it was addressed for delivery to the Commissioner of Patents & Trademarks, Washington, DC 20231 by Express Mail Post Office to Addressee" service.

Name (Print)

Signature

Docket No. 0776/1F216-US2

**IN THE UNITED STATES PATENT AND TRADEMARK OFFICE**

In re Application of:

Allan **SVENDSEN**, *et al.*

Serial No.: To be Assigned  
(Continuation of 08/683,838)

Art Unit:

Filed: Concurrently

Examiner:

For:  **$\alpha$ -AMYLASE MUTANTS**

June 8, 1999

Honorable Commissioner of  
Patents and Trademarks  
Washington, D.C. 20231

**PRELIMINARY AMENDMENT**

Sir:

Prior to examination, applicants wish to amend the above-identified continuation application as follows:

## **IN THE SPECIFICATION**

On page 4, immediately prior to "DETAILED DISCLOSURE OF THE INVENTION"  
please insert:

### **--BRIEF DESCRIPTION OF THE FIGURES**

Fig. 1 shows the 3-dimensional structure of alpha amylase wherein differences in Loop 2 as between Fungamyl-like-alpha amylase and Termamyl-like-alpha amylase are emphasized.

Fig. 2 shows the structure of alpha amylase as in Fig. 1 wherein differences in Loop 3(a) are emphasized.

Fig. 3 shows the structure of alpha amylase as in Fig. 1 wherein differences in Loop 3(b) are emphasized.

Fig. 4 shows the structure of alpha amylase as in Fig. 1 wherein differences in Loop 1(a) are emphasized.

Fig. 5 shows the structure of alpha amylase as in Fig. 1 wherein differences in Loop 1(b) are emphasized.

Fig. 6 shows the structure of the alpha amylase as in Fig. 1 wherein differences in Loop 8 are emphasized.

Fig. 7 shows differences in the alpha amylase structure in the loop between strand 3 and helix 3 of domain A.

Fig. 8 depicts an alpha amylase from WO95/26397.

A map of plasmid pDN1528 is shown in Fig. 9.

A map of plasmid pJeEN1 is shown in Fig. 10.--

**IN THE CLAIMS:**

Please cancel claims 1-69 and add new claims 70-111 as follows:

--70. An alpha-amylase comprising an A domain, a C domain and a calcium binding site, wherein said calcium binding site is associated with said A domain and said C domain comprises ligand residues in said A domain and/or said C domain, wherein said alpha-amylase is modified to alter the characteristics of said calcium binding site and thereby alter the performance of said alpha-amylase by substituting an amino acid residue at a position corresponding to one or more of, Q298, G299, G301, Y302,

L307, N309, Q340, F343, F403, H405, H406, D407, , G410, L427, I428, D430, G433, K436, N473, G474 and G475 in *Bacillus licheniformis*.

--71. The alpha-amylase according to claim 70, wherein said alpha-amylase is produced by *Bacillus*.

--72. The alpha-amylase according to claim 71, wherein said alpha-amylase is produced by *Bacillus licheniformis*, *Bacillus amyloliquefaciens* or *Bacillus stearothermophilus*.

--73. A detergent comprising the alpha-amylase according to claim 70.

--74. A starch liquefaction composition comprising the alpha-amylase according to claim 70.

--75. The alpha-amylase according to claim 70, wherein said alpha-amylase further comprises a substitution or deletion at one or more residues equivalent to M15, V128, H133, W138, N188, A209 and/or M197 in *Bacillus licheniformis*.



--76. The alpha-amylase according to claim 70 which is modified by substituting an amino acid residue at a position corresponding to one or more of G301, H405, H406 and/or K436 in *Bacillus licheniformis*.

--77. An alpha-amylase comprising an A domain, a C domain and a calcium binding site, wherein said calcium binding site is associated with said A domain and said C domain comprises ligand residues in said A domain and/or said C domain, wherein said alpha-amylase is modified to alter the characteristics of said calcium binding site and thereby alter the performance of said alpha-amylase by substituting an amino acid residue at a position corresponding to Q298 in *Bacillus licheniformis*.

--78. An alpha-amylase comprising an A domain, a C domain and a calcium binding site, wherein said calcium binding site is associated with said A domain and said C domain comprises ligand residues in said A domain and/or said C domain, wherein said alpha-amylase is modified to alter the characteristics of said calcium binding site and thereby alter the performance of said alpha-amylase by substituting an amino acid residue at a position corresponding to G299 in *Bacillus licheniformis*.

--79. An alpha-amylase comprising an A domain, a C domain and a calcium binding site, wherein said calcium binding site is associated with said A domain and said C domain comprises ligand residues in said A domain and/or said C domain, wherein

said alpha-amylase is modified to alter the characteristics of said calcium binding site and thereby alter the performance of said alpha-amylase by substituting an amino acid residue at a position corresponding to G301 in *Bacillus licheniformis*.

--80. An alpha-amylase comprising an A domain, a C domain and a calcium binding site, wherein said calcium binding site is associated with said A domain and said C domain comprises ligand residues in said A domain and/or said C domain, wherein said alpha-amylase is modified to alter the characteristics of said calcium binding site and thereby alter the performance of said alpha-amylase by substituting an amino acid residue at a position corresponding to Y302 in *Bacillus licheniformis*.

--81. An alpha-amylase comprising an A domain, a C domain and a calcium binding site, wherein said calcium binding site is associated with said A domain and said C domain comprises ligand residues in said A domain and/or said C domain, wherein said alpha-amylase is modified to alter the characteristics of said calcium binding site and thereby alter the performance of said alpha-amylase by substituting an amino acid residue at a position corresponding to L307 in *Bacillus licheniformis*.

--82. An alpha-amylase comprising an A domain, a C domain and a calcium binding site, wherein said calcium binding site is associated with said A domain and said C domain comprises ligand residues in said A domain and/or said C domain, wherein

said alpha-amylase is modified to alter the characteristics of said calcium binding site and thereby alter the performance of said alpha-amylase by substituting an amino acid residue at a position corresponding to N309 in *Bacillus licheniformis*.

--83. An alpha-amylase comprising an A domain, a C domain and a calcium binding site, wherein said calcium binding site is associated with said A domain and said C domain comprises ligand residues in said A domain and/or said C domain, wherein said alpha-amylase is modified to alter the characteristics of said calcium binding site and thereby alter the performance of said alpha-amylase by substituting an amino acid residue at a position corresponding to Q340 in *Bacillus licheniformis*.

--84. An alpha-amylase comprising an A domain, a C domain and a calcium binding site, wherein said calcium binding site is associated with said A domain and said C domain comprises ligand residues in said A domain and/or said C domain, wherein said alpha-amylase is modified to alter the characteristics of said calcium binding site and thereby alter the performance of said alpha-amylase by substituting an amino acid residue at a position corresponding to F343 in *Bacillus licheniformis*.

--85. An alpha-amylase comprising an A domain, a C domain and a calcium binding site, wherein said calcium binding site is associated with said A domain and said C domain comprises ligand residues in said A domain and/or said C domain, wherein

said alpha-amylase is modified to alter the characteristics of said calcium binding site and thereby alter the performance of said alpha-amylase by substituting an amino acid residue at a position corresponding to F403 in *Bacillus licheniformis*.

--86. An alpha-amylase comprising an A domain, a C domain and a calcium binding site, wherein said calcium binding site is associated with said A domain and said C domain comprises ligand residues in said A domain and/or said C domain, wherein said alpha-amylase is modified to alter the characteristics of said calcium binding site and thereby alter the performance of said alpha-amylase by substituting an amino acid residue at a position corresponding to H405 in *Bacillus licheniformis*.

--87. An alpha-amylase comprising an A domain, a C domain and a calcium binding site, wherein said calcium binding site is associated with said A domain and said C domain comprises ligand residues in said A domain and/or said C domain, wherein said alpha-amylase is modified to alter the characteristics of said calcium binding site and thereby alter the performance of said alpha-amylase by substituting an amino acid residue at a position corresponding to H406 in *Bacillus licheniformis*.

--88. An alpha-amylase comprising an A domain, a C domain and a calcium binding site, wherein said calcium binding site is associated with said A domain and said C domain comprises ligand residues in said A domain and/or said C domain, wherein

said alpha-amylase is modified to alter the characteristics of said calcium binding site and thereby alter the performance of said alpha-amylase by substituting an amino acid residue at a position corresponding to D407 in *Bacillus licheniformis*.

--89. An alpha-amylase comprising an A domain, a C domain and a calcium binding site, wherein said calcium binding site is associated with said A domain and said C domain comprises ligand residues in said A domain and/or said C domain, wherein said alpha-amylase is modified to alter the characteristics of said calcium binding site and thereby alter the performance of said alpha-amylase by substituting an amino acid residue at a position corresponding to G410 in *Bacillus licheniformis*.

--90. An alpha-amylase comprising an A domain, a C domain and a calcium binding site, wherein said calcium binding site is associated with said A domain and said C domain comprises ligand residues in said A domain and/or said C domain, wherein said alpha-amylase is modified to alter the characteristics of said calcium binding site and thereby alter the performance of said alpha-amylase by substituting an amino acid residue at a position corresponding to L427 in *Bacillus licheniformis*.

--91. An alpha-amylase comprising an A domain, a C domain and a calcium binding site, wherein said calcium binding site is associated with said A domain and said C domain comprises ligand residues in said A domain and/or said C domain, wherein

said alpha-amylase is modified to alter the characteristics of said calcium binding site and thereby alter the performance of said alpha-amylase by substituting an amino acid residue at a position corresponding to I428 in *Bacillus licheniformis*.

--92. An alpha-amylase comprising an A domain, a C domain and a calcium binding site, wherein said calcium binding site is associated with said A domain and said C domain comprises ligand residues in said A domain and/or said C domain, wherein said alpha-amylase is modified to alter the characteristics of said calcium binding site and thereby alter the performance of said alpha-amylase by substituting an amino acid residue at a position corresponding to D430 in *Bacillus licheniformis*.

--93. An alpha-amylase comprising an A domain, a C domain and a calcium binding site, wherein said calcium binding site is associated with said A domain and said C domain comprises ligand residues in said A domain and/or said C domain, wherein said alpha-amylase is modified to alter the characteristics of said calcium binding site and thereby alter the performance of said alpha-amylase by substituting an amino acid residue at a position corresponding to G433 in *Bacillus licheniformis*.

--94. An alpha-amylase comprising an A domain, a C domain and a calcium binding site, wherein said calcium binding site is associated with said A domain and said C domain comprises ligand residues in said A domain and/or said C domain, wherein

said alpha -amylase is modified to alter the characteristics of said calcium binding site and thereby alter the performance of said alpha-amylase by substituting an amino acid residue at a position corresponding to K436 in *Bacillus licheniformis*.

--95. An alpha-amylase comprising an A domain, a C domain and a calcium binding site, wherein said calcium binding site is associated with said A domain and said C domain comprises ligand residues in said A domain and/or said C domain, wherein said alpha-amylase is modified to alter the characteristics of said calcium binding site and thereby alter the performance of said alpha-amylase by substituting an amino acid residue at a position corresponding to N473 in *Bacillus licheniformis*.

--96. An alpha-amylase comprising an A domain, a C domain and a calcium binding site, wherein said calcium binding site is associated with said A domain and said C domain comprises ligand residues in said A domain and/or said C domain, wherein said alpha-amylase is modified to alter the characteristics of said calcium binding site and thereby alter the performance of said alpha-amylase by substituting an amino acid residue at a position corresponding to G474 in *Bacillus licheniformis*.

--97. An alpha-amylase comprising an A domain, a C domain and a calcium binding site, wherein said calcium binding site is associated with said A domain and said C domain comprises ligand residues in said A domain and/or said C domain, wherein

said alpha-amylase is modified to alter the characteristics of said calcium binding site and thereby alter the performance of said alpha-amylase by substituting an amino acid residue at a position corresponding to G475 in *Bacillus licheniformis*.

--98. The alpha-amylase of claim 70 wherein said alpha-amylase is modified to alter the characteristics of said calcium binding site and thereby alter the performance of said alpha-amylase by substituting an amino acid residue at a position corresponding to one or more of, Q298, G299, G301, Y302, L307, F343, H405, H406, D407, I428, D430, and G475 in *Bacillus licheniformis*.

--99. The alpha-amylase of claim 98 wherein said alpha-amylase further comprises a substitution or deletion at one or more residues equivalent to A209 and/or M197 in *Bacillus licheniformis*.

--100. The alpha -amylase according to claim 98 which is modified by substituting an amino acid residue at a position corresponding to one or more of G301, H405, and/or H406 in *Bacillus licheniformis*.

--101. The alpha-amylase according to claim 98, wherein said alpha-amylase is produced by *Bacillus*.



--102. The alpha-amylase according to claim 98, wherein said alpha-amylase is produced by *Bacillus licheniformis*, *Bacillus amyloliquefaciens* or *Bacillus stearothermophilus*.

--103. A detergent comprising the alpha-amylase according to claim 98.

--104. A starch liquefaction composition comprising the alpha-amylase according to claim 98.

--105. An alpha-amylase variant of an alpha-amylase having the sequence SEQ. ID # 2 wherein at least one amino acid residue present in a fragment corresponding to amino acid residues 322-346 of SEQ. ID # 2 has been substituted with another residue present in a fragment corresponding to amino acid residues 291-313 of SEQ. ID # 10.

--106. An alpha-amylase wherein at least one amino acid residue present in a fragment corresponding to amino acid residues 322-346 of SEQ. ID # 2 has been substituted with another residue thereby resulting in modified calcium binding properties of said amylase.

--107. An alpha-amylase variant wherein at least one amino acid residue within a distance of 10 Angstrom or less from a calcium ion binding site of said alpha-amylase has been replaced by another amino acid residue thereby modifying the calcium dependence and/or affinity for calcium ion of said amylase.

--108. The alpha-amylase of claim 107 wherein said at least one amino acid residue is selected from the group consisting of V102, I103, N104, H105, K106, R125, W155, W157, Y158, H159, F160, D161, G162, T163, Y175, K176, F177, G178, K180, A181, W182, D183, W184, E185, V186, S187, N192, Y193, D194, Y195, L196, M197, Y198, A199, D200, I201, D202, Y203, D204, H205, P206, V208, A209, D231, A232, V233, K234, H235, I236, K237, F238, F240, L241, A294, A295, S296, T297, Q298, G299, G300, G301, Y302, D303, M304, R305, K306, L307, W342, F343, L346, Q393, Y394, Y396, H405, H406, D407, I408, V409, R413, E414, G415, D416, S417, V419, A420, N421, S422, G423, L424, I428, T429, D430, G431, P432, V440, G441, R442, Q443, N444, A445, G446, E447, T448, W449, I462, G475, Y480, V481, Q482, R483.

--109. The alpha-amylase of claim 108 wherein said at least one amino acid residue that has been replaced is at least two amino acid residues.

--110. An alpha-amylase variant having the sequence SEQ. ID. # 2 wherein at least one of amino acid residues I428-A435 has been substituted in a manner to modify the calcium stability of the alpha-amylase before said substitution.

--111. An alpha-amylase variant having the sequence SEQ. ID. # 2 wherein at least one of amino acid residues T297-L308 has been substituted in a manner to modify the calcium stability of the alpha-amylase before said substitution.

--112. An alpha-amylase variant having the sequence SEQ. ID. # 2 wherein at least one of amino acid residues F403-V409 has been substituted in a manner to improve the calcium stability of the alpha-amylase before said substitution.--

#### **REMARKS**

Consideration of this application is respectfully requested. By the present amendment, the claims of US Patent No. 5,763,385 are copied for purposes of provoking an interference. The amendment to the specification is supported at pp.

8 (lines 30-32); pp. 65 (lines 21-31); pp. 67 (lines 1-2) and in the Figures themselves.

The amendment to the claims is supported *inter alia* at pp. 22-23, 24 and 39-40.

Respectfully submitted,



Adda C. Gogoris  
Registration No. 29,714  
Attorney for Applicants

DARBY & DARBY P.C.  
805 Third Avenue  
New York, New York 10022  
212-527-7700

09/27/2016 10:08:59 AM

## EXPRESS MAIL CERTIFICATE

Date 6/8/99 Label No. 595859379

I hereby certify that, on the date indicated above I deposited this paper or fee with the U.S. Postal Service and that it was addressed for delivery to the Commissioner of Patents & Trademarks, Washington, DC 20237 by "Express Mail Post Office to Addressee" service.

Name (Print) D B PeckSignature D B Peck

PLEASE CHARGE ANY DEFICIENCY UP TO \$300.00  
OR CREDIT ANY EXCESS IN FUTURE FEES DUE  
WITH RESPECT TO THIS APPLICATION TO OUR  
DEPOSIT ACCOUNT NO. 04-0100

Docket No. 0776/1F216-US2

 $\alpha$ -AMYLASE MUTANTS

## 5 FIELD OF THE INVENTION

The present invention relates to a novel method of designing  $\alpha$ -amylase mutants with predetermined properties, which method is based on the hitherto unknown three-dimensional structure of bacterial  $\alpha$ -amylases.

## 10 BACKGROUND OF THE INVENTION

$\alpha$ -Amylases ( $\alpha$ -1,4 glucan-4-glucanohydrolase, EC 3.2.1.1) constitute a group of enzymes which is capable of hydrolyzing starch and other linear and branched 1,4-glucosidic oligo- and polysaccharides. Almost all  $\alpha$ -amylases studied have a few conserved regions with approximately the same length and spacing. One of these regions resembles the  $\text{Ca}^{2+}$  binding site of calmodulin and the others are thought to be necessary for the active centre and/or binding of the substrate.

While the amino acid sequence and thus primary structure of a large number of  $\alpha$ -amylases are known, it has proved very difficult to determine the three-dimensional structure of all  $\alpha$ -amylases. The three-dimensional structure can be determined by X-ray crystallographic analysis of  $\alpha$ -amylase crystals, but it has proven difficult to obtain  $\alpha$ -amylase crystals suitable for actually solving the structure.

Until now the three-dimensional structure of only a few  $\alpha$ -amylases have been determined at high resolution. These include the structure of the *Aspergillus oryzae* TAKA  $\alpha$ -amylase (Swift et al., 1991), the *Aspergillus niger* acid amylase (Brady et al, 1991), the structure of pig pancreatic  $\alpha$ -amylase (Qian et al., 1993), and the barley alpha-amylase (Kadziola et al. 1994, Journal of Molecular Biology 239: 104-121, A.Kadziola, Thesis, Dept of Chemistry, U. of Copenhagen, Denmark). Furthermore, the three-dimensional structure of a *Bacillus circulans* cyclodextrin glycosyltransferase (CGTase) is known (Klein et al., 1992) (Lawson et al., 1994). The CGTase catalyzes the same type of reactions as  $\alpha$ -amylases and exhibits some structural resemblance with  $\alpha$ -amylases.

Furthermore, crystallization and preliminary X-ray studies of *B. subtilis*  $\alpha$ -amylases have been described (Chang et al. (1992) and Mizuno et al. (1993)). No final *B. subtilis* structure has been reported. Analogously, the preparation of *B. licheniformis*

α-amylase crystals has been reported (Suzuki et al. (1990), but no subsequent report on X-ray crystallographic analysis or three-dimensional structure are available.

Several research teams have attempted to build three-dimensional structures on the basis of the above known α-amylase structures. For instance, Vihinen et al. (J. Biochem. 107, 267-272, 1990), disclose the modelling (or computer simulation) of a three-dimensional structure of the *Bacillus stearothermophilus* α-amylase on the basis of the TAKA amylase structure. The model was used to investigate hypothetical structural consequences of various site-directed mutations of the *B. stearothermophilus* α-amylase. E.A. MacGregor (1987) predicts the presence of α-helices and β-barrels in α-amylases from different sources, including barley, pig pancreas and *Bacillus amyloliquefaciens* on the basis of the known structure of the *A. oryzae* TAKA α-amylase and secondary structure predicting algorithms. Furthermore, the possible loops and subsites which may be found to be present in, e.g., the *B. amyloliquefaciens* α-amylase are predicted (based on a comparison with the *A. oryzae* sequence and structure).

A.E. MacGregor (Starch/Stärke 45 (1993), No. 7, p. 232-237) presents a review of the relationship between the structure and activity of α-amylase related enzymes.

Hitherto, no three-dimensional structure has been available for the industrially important *Bacillus* α-amylases (which in the present context are termed "Termamyl-like α-amylases"), including the *B. licheniformis*, the *B. amyloliquefaciens*, and the *B. stearothermophilus* α-amylase.

## BRIEF DISCLOSURE OF THE INVENTION

The three-dimensional structure of a Termamyl-like bacterial α-amylase has now been elucidated. On the basis of an analysis of said structure it is possible to identify structural parts or specific amino acid residues which from structural or functional considerations appear to be important for conferring the various properties to the Termamyl-like α-amylases. Furthermore, when comparing the Termamyl-like α-amylase structure with known structures of the fungal and mammalian α-amylases mentioned above, it has been found that some similarities exist between the structures, but also that some striking, and not previously predicted structural differences between the α-amylases exist. The present invention is based on these findings.

Accordingly, in a first aspect the invention relates to a method of constructing a variant of a parent Termamyl-like  $\alpha$ -amylase, which variant has  $\alpha$ -amylase activity and at least one altered property as compared to said parent  $\alpha$ -amylase, which method comprises

i) analysing the structure of the Termamyl-like  $\alpha$ -amylase with a view to identifying at least one amino acid residue or at least one structural part of the Termamyl-like  $\alpha$ -amylase structure, which amino acid residue or structural part is believed to be of relevance for altering said property of the parent Termamyl-like  $\alpha$ -amylase (as evaluated on the basis of structural or functional considerations),

ii) constructing a Termamyl-like  $\alpha$ -amylase variant, which as compared to the parent Termamyl-like  $\alpha$ -amylase, has been modified in the amino acid residue or structural part identified in i) so as to alter said property, and, optionally,

iii) testing the resulting Termamyl-like  $\alpha$ -amylase variant with respect to said property.

In a second aspect the present invention relates to a method of constructing a variant of a parent Termamyl-like  $\alpha$ -amylase, which variant has  $\alpha$ -amylase activity and one or more altered properties as compared to said parent  $\alpha$ -amylase, which method comprises

i) comparing the three-dimensional structure of the Termamyl-like  $\alpha$ -amylase with the structure of a non-Termamyl-like  $\alpha$ -amylase,

ii) identifying a part of the Termamyl-like  $\alpha$ -amylase structure which is different from the non-Termamyl-like  $\alpha$ -amylase structure,

iii) modifying the part of the Termamyl-like  $\alpha$ -amylase identified in ii) whereby a Termamyl-like  $\alpha$ -amylase variant is obtained, one or more properties of which differ from the parent Termamyl-like  $\alpha$ -amylase, and optionally,

iv) testing the resulting Termamyl-like  $\alpha$ -amylase variant with respect to said property or properties.

In a third aspect the invention relates to a method of constructing a variant of a parent non-Termamyl-like  $\alpha$ -amylase, which variant has  $\alpha$ -amylase activity and one or more altered properties as compared to said parent  $\alpha$ -amylase, which method comprises

i) comparing the three-dimensional structure of the non-Termamyl-like  $\alpha$ -amylase with the structure of a Termamyl-like  $\alpha$ -amylase,

ii) identifying a part of the non-Termamyl-like  $\alpha$ -amylase structure which is different from the Termamyl-like  $\alpha$ -amylase structure,

iii) modifying the part of the non-Termamyl-like  $\alpha$ -amylase identified in ii) whereby a non-Termamyl-like  $\alpha$ -amylase variant is obtained, one or more properties of which differ from the parent non-Termamyl-like  $\alpha$ -amylase, and optionally,

iv) testing the resulting non-Termamyl-like  $\alpha$ -amylase variant with respect to said property or properties.

The property which may be altered by the above methods of the present invention may, e.g., be substrate specificity, substrate binding, substrate cleavage pattern, temperature stability, pH dependent activity, pH dependent stability (especially increased stability at low (e.g. pH<6, in particular pH<5) or high (e.g. pH>9) pH values), stability towards oxidation,  $\text{Ca}^{2+}$ -dependency, specific activity, and other properties of interest. For instance, the alteration may result in a variant which, as compared to the parent Termamyl-like  $\alpha$ -amylase, has an increased specific activity at a given pH and/or an altered substrate specificity.

In still further aspects the invention relates to variants of a Termamyl-like  $\alpha$ -amylase, DNA encoding such variants and methods of preparing the variants. Finally, the invention relates to the use of the variants for various industrial purposes.

## DETAILED DISCLOSURE OF THE INVENTION

### The Termamyl-like $\alpha$ -amylase

It is well known that a number of alpha-amylases produced by *Bacillus* spp. are highly homologous on the amino acid level. For instance, the *B. licheniformis*  $\alpha$ -amylase comprising the amino acid sequence shown in SEQ ID NO: 2 (commercially available as Termamyl®) has been found to be about 89% homologous with the *B. amyloliquefaciens*  $\alpha$ -amylase comprising the amino acid sequence shown in SEQ ID NO: 4 and about 79% homologous with the *B. stearothermophilus*  $\alpha$ -amylase comprising the amino acid sequence shown in SEQ ID NO: 6. Further homologous  $\alpha$ -amylases include an  $\alpha$ -amylase derived from a strain of the *Bacillus* sp. NCIB 12289, NCIB 12512, NCIB 12513 or DSM 9375, all of which are described in detail in WO 95/26397, and the  $\alpha$ -amylase described by Tsukamoto et al., 1988, Biochemical and Biophysical Research Communications, Vol. 151, No. 1. Still other homologous  $\alpha$ -amylases include the  $\alpha$ -amylase produced by the *B. licheniformis* described in EP 252 666 (ATCC 27811), and the  $\alpha$ -amylases identified in WO 91/00353 and WO 94/18314. Other commercial Termamyl-like *B. licheniformis*  $\alpha$ -amylases are Optitherm®



and Takatherm® (available from Solvay), Maxamyl® (available from Gist-brocades/Genencor), Spezym AA® (available from Genencor), and Keistase® (available from Daiwa).

Because of the substantial homology found between these  $\alpha$ -amylases, they are considered to belong to the same class of  $\alpha$ -amylases, namely the class of "Termamyl-like  $\alpha$ -amylases".

Accordingly, in the present context, the term "Termamyl-like  $\alpha$ -amylase" is intended to indicate an  $\alpha$ -amylase which, on the amino acid level, exhibits a substantial homology to Termamyl®, i.e. the *B. licheniformis*  $\alpha$ -amylase SEQ ID NO: 2. In other words, a Termamyl-like  $\alpha$ -amylase is an  $\alpha$ -amylase, which has the amino acid sequence shown in SEQ ID NO: 2, 4 or 6 herein, or the amino acid sequence shown in SEQ ID NO: 1 or 2 of WO 95/26397 or in Tsukamoto et al., 1988, or i) which displays at least 60%, such as at least 70%, e.g. at least 75%, or at least 80%, e.g. at least 85%, at least 90% or at least 95% homology with at least one of said amino acid sequences and/or ii) displays immunological cross-reactivity with an antibody raised against at least one of said  $\alpha$ -amylases, and/or iii) is encoded by a DNA sequence which hybridizes to the DNA sequences encoding the above specified  $\alpha$ -amylases which are apparent from SEQ ID NO: 1, 3 and 5 of the present application, and SEQ ID NO: 4 and 5 of WO 95/26397, respectively.

In connection with property i) the "homology" may be determined by use of any conventional algorithm, preferably by use of the GAP programme from the GCG package version 7.3 (June 1993) using default values for GAP penalties (Genetic Computer Group (1991) Programme Manual for the GCG Package, version 7, 575 Science Drive, Madison, Wisconsin, USA 53711).

Property ii) of the  $\alpha$ -amylase, i.e. the immunological cross reactivity, may be assayed using an antibody raised against or reactive with at least one epitope of the relevant Termamyl-like  $\alpha$ -amylase. The antibody, which may either be monoclonal or polyclonal, may be produced by methods known in the art, e.g. as described by Hudson et al., 1989. The immunological cross-reactivity may be determined using assays known in the art, examples of which are Western Blotting or radial immunodiffusion assay, e.g. as described by Hudson et al., 1989. In this respect, immunological cross-reactivity between the  $\alpha$ -amylases having the amino acid sequences SEQ ID NO: 2, 4 and 6, respectively, has been found.

The oligonucleotide probe used in the characterization of the Termamyl-like  $\alpha$ -amylase in accordance with property iii) above may suitably be prepared on the basis of the full or partial nucleotide or amino acid sequence of the  $\alpha$ -amylase in question. Suitable conditions for testing hybridization involve presoaking in 5xSSC and prehybridizing for 1h at  $\sim 40^{\circ}\text{C}$  in a solution of 20% formamide, 5xDenhardt's solution, 50mM sodium phosphate, pH 6.8, and 50 $\mu\text{g}$  of denatured sonicated calf thymus DNA, followed by hybridization in the same solution supplemented with 100 $\mu\text{M}$  ATP for 18h at  $\sim 40^{\circ}\text{C}$ , or other methods described by e.g. Sambrook et al., 1989.

In the present context, "derived from" is intended not only to indicate an  $\alpha$ -amylase produced or producible by a strain of the organism in question, but also an  $\alpha$ -amylase encoded by a DNA sequence isolated from such strain and produced in a host organism transformed with said DNA sequence. Finally, the term is intended to indicate an  $\alpha$ -amylase which is encoded by a DNA sequence of synthetic and/or cDNA origin and which has the identifying characteristics of the  $\alpha$ -amylase in question. The term is also intended to indicate that the parent  $\alpha$ -amylase may be a variant of a naturally occurring  $\alpha$ -amylase, i.e. a variant which is the result of a modification (insertion, substitution, deletion) of one or more amino acid residues of the naturally occurring  $\alpha$ -amylase.

#### Parent hybrid $\alpha$ -amylases

The parent  $\alpha$ -amylase (being a Termamyl-like or non-Termamyl-like  $\alpha$ -amylase) may be a hybrid  $\alpha$ -amylase, i.e. an  $\alpha$ -amylase which comprises a combination of partial amino acid sequences derived from at least two  $\alpha$ -amylases.

The parent hybrid  $\alpha$ -amylase may be one which on the basis of amino acid homology and/or immunological cross-reactivity and/or DNA hybridization (as defined above) can be determined to belong to the Termamyl-like  $\alpha$ -amylase family. In this case, the hybrid  $\alpha$ -amylase is typically composed of at least one part of a Termamyl-like  $\alpha$ -amylase and part(s) of one or more other  $\alpha$ -amylases selected from Termamyl-like  $\alpha$ -amylases or non-Termamyl-like  $\alpha$ -amylases of microbial (bacterial or fungal) and/or mammalian origin.

Thus, the parent hybrid  $\alpha$ -amylase may comprise a combination of at least two Termamyl-like  $\alpha$ -amylases, or of at least one Termamyl-like and at least one non-Termamyl-like bacterial  $\alpha$ -amylase, or of at least one Termamyl-like and at least one fungal  $\alpha$ -amylase. For instance, the parent  $\alpha$ -amylase comprises a C-terminal part of an  $\alpha$ -amylase derived from

a strain of *B. licheniformis* and a N-terminal part of an  $\alpha$ -amylase derived from a strain of *B. amyloliquefaciens* or from a strain of *B. stearothermophilus*. For instance, the parent  $\alpha$ -amylase comprises at least 430 amino acid residues of the C-terminal part of the *B. licheniformis*  $\alpha$ -amylase, and may, e.g. comprise a) an amino acid segment corresponding to the 37 N-terminal amino acid residues of the *B. amyloliquefaciens*  $\alpha$ -amylase having the amino acid sequence shown in SEQ ID NO: 4 and an amino acid segment corresponding to the 445 C-terminal amino acid residues of the *B. licheniformis*  $\alpha$ -amylase having the amino acid sequence shown in SEQ ID NO: 2, or b) an amino acid segment corresponding to the 68 N-terminal amino acid residues of the *B. stearothermophilus*  $\alpha$ -amylase having the amino acid sequence shown in SEQ ID NO: 6 and an amino acid segment corresponding to the 415 C-terminal amino acid residues of the *B. licheniformis*  $\alpha$ -amylase having the amino acid sequence shown in SEQ ID NO: 2.

Analogously, the parent hybrid  $\alpha$ -amylase may belong to a non-Termamyl-like  $\alpha$ -amylase family, e.g. the Fungamyl-like  $\alpha$ -amylase family. In that case the hybrid may comprise at least one part of an  $\alpha$ -amylase belonging to the non-Termamyl-like  $\alpha$ -amylase family in combination with one or more parts derived from other  $\alpha$ -amylases.

#### The three-dimensional Termamyl-like $\alpha$ -amylase structure

The Termamyl-like  $\alpha$ -amylase which was used to elucidate the three-dimensional structure forming the basis for the present invention consists of the 300 N-terminal amino acids of the *B. amyloliquefaciens*  $\alpha$ -amylase (with the amino acid sequence shown in SEQ ID NO: 4) and amino acids 301-483 of the C-terminal end of the *B. licheniformis*  $\alpha$ -amylase with the amino acid sequence SEQ ID NO: 2. The bacterial  $\alpha$ -amylase belongs to the "Termamyl-like  $\alpha$ -amylase family" and the present structure is believed to be representative for the structure of any Termamyl-like  $\alpha$ -amylase.

The structure of the  $\alpha$ -amylase was solved in accordance with the principle for X-ray crystallographic methods given in "X-Ray Structure Determination", Stout, G.K. and Jensen, L.H., John Wiley & Sons, inc. NY, 1989. The structural coordinates for the solved crystal structure of the  $\alpha$ -amylase at 2.2 Å resolution using the isomorphous replacement method are given in a standard PDB format (Brookhaven Protein Data Base) in Appendix 1. It is to be understood that Appendix 1 forms part of the present application.

Amino acid residues of the enzyme are identified by three-letter amino acid code (capitalized letters).

The  $\alpha$ -amylase structure is made up of three globular domains ordered A, B, and C with respect to sequence, which lie approximately along a line in the order B, A, C. The domains can be defined as being residues 1-103 and 206-395 for domain A, residues 104-205 for domain B, and residues 396-483 for domain C, the numbers referring to the *B. licheniformis*  $\alpha$ -amylase. This gives rise to an elongated molecule, the longest axis being about 85Å. The widest point perpendicular to this axis is approximately 50Å and spans the central A domain. The active site residues of the *B. licheniformis*  $\alpha$ -amylase (SEQ ID NO: 2) are D323, D231 and E261.

#### Domain A

Domain A is the largest domain and contains the active site (comprised of a cluster of three amino acid residues placed at the bottom of a deep cleft in the enzyme's surface). Domain A of all known  $\alpha$ -amylase structures have the same overall fold, viz. the (beta/alpha)<sub>8</sub> barrel with 8 central beta strands (number 1-8) and 8 flanking  $\alpha$ -helices. The  $\beta$ -barrel is defined by McGregor *op. cit.* The C-terminal end of Beta strand 1 is connected to helix 1 by a loop denoted loop 1 and an identical pattern is found for the other loops. These loops show some variation in size and some can be quite extensive.

The 8 central Beta-strands in the (beta/alpha)<sub>8</sub> barrel superimpose well between the various known  $\alpha$ -amylase structures, and this part of the structure, including the close surroundings of the active site located at the c-terminal end of the beta-strands, show high similarity between the different amylases.

The loops connecting beta-strands and alpha helices display high variations between alpha amylases. These loops constitute the structural context of the active site and the majority of the contacts to the substrate is found among residues located in these loops. Such important characteristics as substrate specificity, substrate binding, pH/activity profile, starch cleavage pattern are determined by the amino acids and the positions of same in these loops.

The substantial differences between the Fungamyl-like  $\alpha$ -amylase structure and the structure of the Termamyl-like  $\alpha$ -amylase disclosed herein which are found in loops 1, 2, 3, and 8 are visualized in the Figures.

### Domain B

The Termamyl-like  $\alpha$ -amylase structure has been found to comprise a special domain structure in the A domain's loop3, also called domain B. The structure of the Termamyl-like  $\alpha$ -amylase B domain has never been seen before in any of the known  $\alpha$ -amylase or ( $\beta$ /alpha)8-barrel proteins.

The domain B structure is a very compact domain having a very high number of charged residues. The B domain arises as an extension of the loop between strand 3 and helix 3 of domain A (shown in Fig. 7) and contains a 5 stranded antiparallel  $\beta$ -sheet structure containing at least one long loop structure and having the connectivity -1, +3, -1X, +2 (Richardson, 1981, Adv. Protein Chem. 34, 167-339).

The first four strands of the B domain form two hairpin loops which twist around each other like a pair of crossed fingers (right-hand twist). The mainchain folds into a  $\beta$ -strand which connects two small  $\beta$ -sheet structures. After making one turn in one sheet it folds back and makes up a two stranded sheet in contact with domain A and an internal hole in the  $\alpha$ -amylase structure. Then the mainchain folds up to a small sheet structure nearly perpendicular to the first two sheets. Before entering the helix 3 on top of the  $\beta$ -strand 3, the approximately 24 last amino acids in domain B form two calcium binding sites in the contact region to domain A.

Domain B is connected with domain A by two peptide stretches, which divide the domain-domain contact areas into two. Domain B is in contact with Domain A by a calcium binding region and an internally buried hole containing waters. Many types of molecular contacts are present. Ionic interacting between acid and basic amino acids are possible, these interactions are very important for the general stability at high pH and for keeping the Calcium binding sites intact.

### Domain C

Domain C is the C-terminal part of the protein consisting of amino acids 394-483. Domain C is composed entirely of  $\beta$ -strands which forms a single 8-stranded sheet structure, which folds back on itself, and thus may be described as a  $\beta$ -sandwich structure. The connectivity is +1,+1, +5, -3, +1, +1, -3 although strands 6 and 7 are only loosely connected. One part of the  $\beta$ -sheet forms the interface to domain A.

### Ca-binding and Na-binding sites

The structure of the Termamyl-like  $\alpha$ -amylase is remarkable in that it exhibits four calcium-binding sites and one sodium-binding site. In other words four calcium ions and one sodium ion are found to be present in the structure, although one of the calcium ions displays very weak coordination. Two of the calcium ions form part of a linear cluster of three ions, the central ion being attributed to sodium, which lie at the junction of the A and B domains.

The coordinating residues for the calcium ions between the A and B domain are as follows (using the Pdb file nomenclature for amino acid residues and atoms in the Pdb file found in Appendix 1 herein): For the calcium ion nearest to the active site (IUM 502 in the pdb file), the backbone carbonyls from His235 and Asp194, the sidechain atom OD1 from residues Asp194, Asn102 and Asp200, and one water molecule WAT X3 (atom OW7). For the sodium ion (IUM 505), the binding site includes atom OD2 from Asp194, Asp200, Asp183 and Asp159, and a backbone carbonyl from Val201. The coordinates for the other calcium ion between domain A and B are (IUM 501) : atom OD2 from Asp204 and Asp159, backbone carbonyl from Asp183 and Ala181, atom OD1 from Asp202, and one water molecule WAT X7 (atom OW7).

One calcium ion is located between the A and C domain, another is located in the C domain. The first mentioned calcium ion, which is also the one best coordinated (IUM 503) includes a carbonyl backbone from Gly300, Tyr302 and His406, atom OD2/OD1 from Asp430, atom OD1 from Asp407, and one water molecule WAT X6 (atom OW7). The other and very weakly coordinated calcium site (IUM 504) comprises 4 water molecules WAT X21 (atom OW8), X6 (atom OW6), X9 (atom OW0) and X28 (atom OW8), OE1/OE2 from Glu447 and OD1 from Asn444.

### Substrate-binding site

Without being limited to any theory it is presently believed that favourable interactions between a substrate molecule and the enzyme (such as hydrogen bonds and/or strong electrostatic interaction) are found within a sphere of 4Å of the substrate, when bound to the enzyme. The following residues of the *B. licheniformis*  $\alpha$ -amylase having the amino acid sequence shown in SEQ ID NO: 2 are contemplated to be within a distance of 4 Å of the substrate and thus believed to be involved in interactions with the substrate:

Trp13, Tyr14, Asn17, Asp18, Ser50, Gln51, Ala52, Asp53, Val54, Gly55, Tyr56, Lys70, Arg74, Lys76, Val102, His105, Gly107, Gly108, Ala109, Trp138, Thr163, Asp164, Trp165, Asn172, Glu189, Tyr193, Leu196, Met197, Tyr198, Ala199, Arg229, Asp231, Ala232, Lys234, His235, Glu261, Trp263, His327, Asp328, Gln333, Ser334, and Leu335.

5 The amino acid residues of another Termamyl-like  $\alpha$ -amylase, which are contemplated to be within a distance of 4Å of the substrate, may easily be identified by aligning the amino acid sequence SEQ ID NO: 2 with that of the other Termamyl-like  $\alpha$ -amylase and thereby identifying the positions equivalent to those identified above.

#### 10 Generality of structure

Because of the high homology between the various Termamyl-like  $\alpha$ -amylases, the solved structure defined by the coordinates of Appendix 1 is believed to be representative for the structure of all Termamyl-like  $\alpha$ -amylases. A model structure of other Termamyl-like  $\alpha$ -amylases may easily be built on the basis of the coordinates given in Appendix 1 adapted to the  $\alpha$ -amylase in question by use of an alignment between the respective amino acid sequences. The creation of a model structure is exemplified in Example 1.

15 The above identified structurally characteristic parts of the Termamyl-like  $\alpha$ -amylase structure (Ca-binding site, substrate binding site, loops, etc.) may easily be identified in other Termamyl-like  $\alpha$ -amylases on the basis of a model (or solved) structure of the relevant Termamyl-like  $\alpha$ -amylase or simply on the basis of an alignment between the amino acid sequence of the Termamyl-like  $\alpha$ -amylase in question with that of the *B. lichenformis*  $\alpha$ -amylase used herein for identifying the amino acid residues of the respective structural elements.

20 Furthermore, in connection with Termamyl-like variants of the invention, which are defined by modification of specific amino acid residues of a specific Termamyl-like  $\alpha$ -amylase, it will be understood that variants of another Termamyl-like  $\alpha$ -amylase modified in an equivalent position (as determined from the best possible amino acid sequence alignment between the respective sequences) are intended to be covered as well. Thus, irrespective of whether an amino acid residue is identified herein for the purpose of defining a structural part of a given  $\alpha$ -amylase or used for identifying a variant of the  $\alpha$ -amylase, this amino acid residue shall be considered as representing the equivalent amino acid residue of any other Termamyl-like  $\alpha$ -amylase.

25

30

### Methods of the invention for design of novel $\alpha$ -amylase variants

In the methods according to the first, second and third aspects of the invention the terms "structure of a Termamyl-like  $\alpha$ -amylase" and "Termamyl-like  $\alpha$ -amylase structure" are intended to indicate the solved structure defined by the coordinates presented in Appendix 1 or a model structure of a given Termamyl-like  $\alpha$ -amylase (such as the *B. licheniformis*  $\alpha$ -amylase) built on the basis of the solved structure.

In most cases the parent Termamyl-like  $\alpha$ -amylase to be modified in accordance with the present invention is different from the  $\alpha$ -amylase which was actually used for solving the structure (Appendix 1). This means that the amino acid residue(s) or structural part(s) identified in the solved structure (Appendix 1) in step i) of the method according to the first, second or third aspect of the invention must be translated into the corresponding amino acid residue(s) or structural part(s) of the parent Termamyl-like  $\alpha$ -amylase in question. The "translation" is conveniently performed on the basis of an amino acid sequence alignment between the amino acid sequence of the Termamyl-like  $\alpha$ -amylase used for solving the structure and the amino acid sequence of the parent Termamyl-like  $\alpha$ -amylase in question.

The analysis or comparison performed in step i) of the method according to the first, second and third aspect, respectively, of the invention may be performed by use of any suitable computer programme capable of analysing and/or comparing protein structures, e.g. the computer programme Insight, available from Biosym Technologies, Inc. For instance, the basic principle of structure comparison is that the three-dimensional structures to be compared are superimposed on the basis of an alignment of secondary structure elements (such as the central 8  $\beta$ -strands in the barrel) and the parts differing between the structures can subsequently easily be identified from the superimposed structure.

The structural part which is identified in step i) of the methods of the first, second and third aspects of the invention may be composed of one amino acid residue. However, normally the structural part comprises more than one amino acid residue, typically constituting one of the above parts of the Termamyl-like  $\alpha$ -amylase structure such as one of the A, B, or C domains, an interface between any of these domains, a calcium binding site, a loop structure, the substrate binding site, or the like.

In the present context the term "structural or functional considerations" is intended to indicate that modifications are made on the basis of an analysis of the relevant structure or structural part and its contemplated impact on the function of the enzyme. Thus,



an analysis of the structures of the various  $\alpha$ -amylases, which until now has been elucidated, optionally in combination with an analysis of the functional differences between these  $\alpha$ -amylases, may be used for assigning certain properties of the  $\alpha$ -amylases to certain parts of the  $\alpha$ -amylase structure or to contemplate such relationship. For instance, differences in the pattern or structure of loops surrounding the active site may result in differences in access to the active site of the substrate and thus differences in substrate specificity and/or cleavage pattern. Furthermore, parts of a Termamyl-like  $\alpha$ -amylase involved in or contemplated to be involved in substrate binding (and thus e.g. specificity/cleavage pattern), calcium or sodium ion binding (e.g. of importance for the Calcium-dependency of the enzyme), and the like has been identified (*vide infra*).

The modification of an amino acid residue or structural part is typically accomplished by suitable modifications of a DNA sequence encoding the parent enzyme in question. The term "modified" as used in step ii) in the method according to the first aspect of the invention is intended to have the following meaning: When used in relation to an amino acid residue the term is intended to mean replacement of the amino acid residue in question with another amino acid residue. When used in relation to a structural part, the term is intended to mean replacement of one or more amino acid residues of said structural part, addition of one or more amino acid residues to said part, or deletion of one or more amino acid residues of said structural part.

The construction of the variant of interest is accomplished by cultivating a microorganism comprising a DNA sequence encoding the variant under conditions which are conducive for producing the variant, and optionally subsequently recovering the variant from the resulting culture broth. This is described in detail further below.

#### *First aspect of the invention*

In a preferred embodiment of the method according to the first aspect of the invention the property of the parent enzyme to be modified is selected from calcium dependency, substrate binding, cleavage pattern, pH dependent activity and the like. Specific examples of how to change these properties of a parent Termamyl-like  $\alpha$ -amylase are given further below.

In another preferred embodiment the parent Termamyl-like  $\alpha$ -amylase to be modified is a *B. licheniformis*  $\alpha$ -amylase.

## *Second and third aspects of the invention*

One important advantage of the methods according to the second and third aspects of the present invention is that it is possible to adapt the structure (or a structural part) of a Termamyl-like  $\alpha$ -amylase to the structure (or structural part) of a non-Termamyl-like  $\alpha$ -amylase and *vice versa*. For instance, having identified a loop structure of the non-Termamyl-like  $\alpha$ -amylase which is believed to be responsible for or contributing to a particular property of the non-Termamyl-like  $\alpha$ -amylase it is possible to replace the corresponding structure of the Termamyl-like  $\alpha$ -amylase with said non-Termamyl-like  $\alpha$ -amylase structure - or if no corresponding structure exists in the Termamyl-like  $\alpha$ -amylase - to insert the structure into the Termamyl-like  $\alpha$ -amylase in such a manner that the resulting variant Termamyl-like  $\alpha$ -amylase, as far as the relevant part is concerned, resembles the corresponding part of the non-Termamyl-like  $\alpha$ -amylase. When two or more parts of the structure of the parent Termamyl-like  $\alpha$ -amylase are modified so as to resemble the corresponding parts of the non-Termamyl-like  $\alpha$ -amylase it is possible to increase the resemblance to the non-Termamyl-like  $\alpha$ -amylase of the Termamyl-like  $\alpha$ -amylase variant and thus to alter the properties of said variant in the direction of those of said non-Termamyl-like  $\alpha$ -amylase. Loop modifications are discussed in much further detail further below.

Typically, the modification to be performed in step iii) of the method according to the second aspect of the invention is accomplished by deleting one or more amino acid residues of the part of the Termamyl-like  $\alpha$ -amylase to be modified so as to adapt the structure of said part of the parent  $\alpha$ -amylase to the corresponding part of the non-Termamyl-like  $\alpha$ -amylase; by replacing one or more amino acid residues of the part of the Termamyl-like  $\alpha$ -amylase to be modified with the amino acid residues occupying corresponding positions in the non-Termamyl-like  $\alpha$ -amylase; or by insertion of one or more amino acid residues present in the non-Termamyl-like  $\alpha$ -amylase into a corresponding position in the Termamyl-like  $\alpha$ -amylase. For the method according to the third aspect the modification is to be understood analogously, performed on the non-Termamyl-like parent  $\alpha$ -amylase rather than the Termamyl-like  $\alpha$ -amylase.

In step ii) of the method according to the second or third aspect of the invention the part of the structure to be identified is preferably one which in the folded enzyme is believed to be in contact with the substrate (cf the disclosure above in the section entitled "Substrate-binding site) or involved in substrate specificity and/or cleavage pattern.

and/or one which is in contact with one of the calcium or sodium ions and/or one, which is contributing to the pH or temperature profile of the enzyme, or one which otherwise, from structural or functional considerations, is contemplated to be responsible for differences in one or more properties of the Termamyl-like and non-Termamyl-like  $\alpha$ -amylase.

#### Non-Termamyl-like $\alpha$ -amylase

The non-Termamyl-like  $\alpha$ -amylase with which the comparison is made in step i) of the method of the second aspect of the invention and which is the parent  $\alpha$ -amylase in the method of the third aspect of the invention, may be any  $\alpha$ -amylase, which does not belong to the family of Termamyl-like  $\alpha$ -amylases (as defined above) and, which as a consequence thereof, has a different three-dimensional structure. Furthermore, the non-Termamyl-like  $\alpha$ -amylase should be one which has, at the time that the method is performed, an elucidated or contemplated three-dimensional structure.

The non-Termamyl-like  $\alpha$ -amylase may, e.g., be a fungal  $\alpha$ -amylase, a mammalian or a plant  $\alpha$ -amylase or a bacterial  $\alpha$ -amylase (different from a Termamyl-like  $\alpha$ -amylase). Specific examples of such  $\alpha$ -amylases include the *Aspergillus oryzae* TAKA  $\alpha$ -amylase, the *A. niger* acid  $\alpha$ -amylase, the *Bacillus subtilis*  $\alpha$ -amylase, the porcine pancreatic  $\alpha$ -amylase and a barley  $\alpha$ -amylase. All of these  $\alpha$ -amylases have elucidated structures which are clearly different from the structure of the Termamyl-like  $\alpha$ -amylase shown herein.

The fungal  $\alpha$ -amylases mentioned above, i.e. derived from *A. niger* and *A. oryzae*, are highly homologous on the amino acid level and generally considered to belong to the same family of  $\alpha$ -amylases. In the present disclosure, this family is termed "Fungamyl-like  $\alpha$ -amylase" and intends to indicate an  $\alpha$ -amylase which exhibits a high homology, i.e. more than 70%, such as 80% homologous (as defined herein) to the fungal  $\alpha$ -amylase derived from *Aspergillus oryzae*, commercially available as Fungamyl®, and the *A. niger*  $\alpha$ -amylase.

From the enclosed illustrations of the  $\alpha$ -amylase structure of a Termamyl-like  $\alpha$ -amylase and a comparison of said structure with the structure of a Fungamyl-like  $\alpha$ -amylase it is evident that major differences exist between the two structures. In the method of the invention it is of particular interest to modify parts of the parent Termamyl-like  $\alpha$ -amylase, which belong to a region with large differences to the Fungamyl-like  $\alpha$ -amylase. In particular, it is of interest to modify the parent Termamyl-like  $\alpha$ -amylase in one or more of the following loops: loop 1, loop 2, loop 3 and/or loop 8 of the parent  $\alpha$ -amylase.

In the method of the third aspect of the invention it is of particular interest to modify loop 1, loop 2, loop 3 and/or loop 8 of the parent non-Termamyl-like  $\alpha$ -amylase to a closer resemblance to the similar loops of a Termamyl-like  $\alpha$ -amylase, such as Termamyl.

In the following specific types of variants are described which have been designed by use of the method of the invention.

#### Loop modifications

In order to change the substrate specificity of the parent  $\alpha$ -amylase to be modified it is relevant to consider loop modifications. For instance changing one or more of the loop structures of the Termamyl-like  $\alpha$ -amylase into a closer resemblance with the corresponding loop structure(s) of a non-Termamyl-like  $\alpha$ -amylase (such as a Fungamyl-like  $\alpha$ -amylase) it is contemplated that it is possible to change the substrate specificity in the direction of that of the non-Termamyl  $\alpha$ -amylase. In the following different types of loop modifications of interest are listed. It will be understood that the variants may have other changed properties in addition to the modified substrate specificity. It will be understood that the following modifications identified for a specific Termamyl-like  $\alpha$ -amylase are intended to include corresponding modifications in other equivalent positions of other Termamyl-like  $\alpha$ -amylases. Furthermore, it will be understood that, normally, the loop modification will comprise replacement of an entire loop structure or a substantial part thereof in, e.g., the Termamyl-like  $\alpha$ -amylase, with the corresponding loop structure (or substantial part thereof) in a non-Termamyl-like  $\alpha$ -amylase.

#### *Loop2 modifications*

In one embodiment the invention relates to a variant of a parent Termamyl-like  $\alpha$ -amylase, in which variant at least one amino acid residue of the parent  $\alpha$ -amylase, which is/are present in a fragment corresponding to the amino acid fragment 44-57 of the amino acid sequence of SEQ ID NO: 4, i.e. loop 2, has been deleted or replaced with one or more amino acid residues which is/are present in a fragment corresponding to the amino acid fragment 66-84 of the amino acid sequence shown in SEQ ID NO: 10, or in which one or more additional amino acid residues has been added using the relevant part of SEQ ID NO: 10 or a corresponding part of another Fungamyl-like  $\alpha$ -amylase as a template.

The amino acid sequence shown in SEQ ID NO: 10 is the amino acid sequence of the *A. oryzae*  $\alpha$ -amylase, i.e. a Fungamyl-like  $\alpha$ -amylase. It will be understood that amino acid residues or fragments found in corresponding positions in other  $\alpha$ -amylases, in particular Fungamyl-like  $\alpha$ -amylases, may be used as a template for the construction of the variant according to the invention. The corresponding part in other homologous  $\alpha$ -amylases may easily be identified on the basis of a comparison of the amino acid sequences and/or three-dimensional structures of the respective  $\alpha$ -amylases.

For instance, the variant may be one, which, when the amino acid sequence of the variant is aligned most closely with the amino acid sequence of the said parent  $\alpha$ -amylase, occupies the same position as the portion from residue X to residue Y of SEQ ID NO: 4, the said region having at least 80% such as at least 90% sequence homology with the part of SEQ ID NO: 10 extending from residue Z to residue V of SEQ ID NO: 10, wherein

X is the amino acid residue occupying position 44, 45, 46, 47 or 48 of SEQ ID NO: 4,

Y is the amino acid residue occupying position 51, 52, 53, 54, 55, 56 or 57 of SEQ ID NO: 4,

Z is the amino acid residue occupying position 66, 67, 68, 69 or 70 of SEQ ID NO: 10, and

V is the amino acid residue occupying position 78, 79, 80, 81, 82, 83 or 84 of SEQ ID NO: 10.

In other words, the variant may be one in which an amino acid fragment X-Y of the parent  $\alpha$ -amylase, which corresponds to or is within the amino acid fragment 44-57 of SEQ ID NO: 4, has been replaced with an amino acid fragment Z-V, which corresponds to or is within the amino acid fragment 66-84 of the amino acid sequence shown in SEQ ID NO: 10, in X, Y, Z and V have the meaning indicated above.

A specific example of a variant according to this embodiment is a variant of a parent Termamyl-like  $\alpha$ -amylase, in which the amino acid fragment of the parent  $\alpha$ -amylase, which corresponds to amino acid residues 48-51 of SEQ ID NO: 4, has been replaced with an amino acid fragment corresponding to amino acid residues 70-78 of the amino acid sequence shown in SEQ ID NO: 10.

### *Loop 3 modifications - limited alteration*

In another embodiment the invention relates to a variant of a parent Termamyl-like  $\alpha$ -amylase, in which variant at least one of the amino acid residues of the parent  $\alpha$ -amylase, which is/are present in an amino acid fragment corresponding to the amino acid fragment 195-202 of the amino acid sequence of SEQ ID NO: 4, has been deleted or replaced with one or more of the amino acid residues which is/are present in an amino acid fragment corresponding to the amino acid fragment 165-177 of the amino acid sequence shown in SEQ ID NO: 10, or in which one or more additional amino acid residues has been added using the relevant part of SEQ ID NO: 10 or a corresponding part of another Fungamyl-like  $\alpha$ -amylase as a template.

For instance, the variant may be one in which an amino acid fragment X-Y of the parent  $\alpha$ -amylase which corresponds to or is within the amino acid fragment 195-202 of SEQ ID NO: 4, has been replaced by an amino acid fragment Z-V, which corresponds to or is within the amino acid fragment 165-177 of the amino acid sequence shown in SEQ ID NO: 10, in which

X is an amino acid residue corresponding to the amino acid occupying position 195 or 196 of SEQ ID NO: 4,

Y is an amino acid residue corresponding to the amino acid occupying position 198, 199, 200, 201, or 202 of SEQ ID NO: 4,

Z is an amino acid residue corresponding to the amino acid occupying position 165 or 166 of SEQ ID NO: 10, and

V is an amino acid residue corresponding to the amino acid occupying position 173, 174, 175, 176 or 177 of SEQ ID NO: 10.

Expressed in another manner, the variant according to this aspect may be one, which, when the amino acid sequence of variant is aligned most closely with the amino acid sequence of the said parent Termamyl-like  $\alpha$ -amylase, occupies the same position as the portion from residue X to residue Y of SEQ ID NO: 4, the said region having at least 80%, such as 90% sequence homology with the part of SEQ ID NO: 10 extending from residue Z to residue V of SEQ ID NO: 10, the meaning of X, Y, Z and V being as identified above.

A specific example of a variant according to this embodiment is a variant of a parent Termamyl-like  $\alpha$ -amylase, in which the amino acid fragment of the parent  $\alpha$ -amylase, which corresponds to amino acid residues 196-198 of SEQ ID NO: 4, has been replaced with

the amino acid fragment corresponding to amino acid residues 166-173 of the amino acid sequence shown in SEQ ID NO: 10.

*Loop 3 modifications - complete domain B*

5 In a further embodiment the invention relates to a variant of a parent Termamyl-like  $\alpha$ -amylase, in which variant at least one of the amino acid residues of the parent  $\alpha$ -amylase, which is/are present in a fragment corresponding to the amino acid fragment 117-185 of the amino acid sequence of SEQ ID NO: 4, has/have been deleted or replaced with one or more of the amino acid residues, which is/are present in an amino acid  
10 fragment corresponding to the amino acid fragment 98-210 of the amino acid sequence shown in SEQ ID NO: 10, or in which one or more additional amino acid residues has been added using the relevant part of SEQ ID NO: 10 or a corresponding part of another Fungamyl-like  $\alpha$ -amylase as a template.

For instance, the variant may be one, in which an amino acid fragment X-Y  
15 of the parent  $\alpha$ -amylase, which corresponds to or is within the amino acid fragment 117-185 of SEQ ID NO: 4, has been replaced with an amino acid fragment Z-V, which corresponds to or is within the amino acid fragment 98-210 of the amino acid sequence shown in SEQ ID NO: 10, in which variant

X is an amino acid residue corresponding to the amino acid occupying position  
20 117, 118, 119, 120 or 121 of SEQ ID NO: 4,

Y is an amino acid residue corresponding to the amino acid occupying position  
181, 182, 183, 184 or 185 of SEQ ID NO: 4,

Z is an amino acid residue corresponding to the amino acid occupying position  
98, 99, 100, 101, 102 of SEQ ID NO: 10, and

25 V is an amino acid residue corresponding to the amino acid occupying position 206, 207, 208, 209 or 210 of SEQ ID NO: 10.

A specific example of a variant according to this embodiment is a variant of a parent  $\alpha$ -amylase, in which an amino acid fragment of the parent  $\alpha$ -amylase, which corresponds to amino acid residues 121-181 of SEQ ID NO: 4, has been replaced with the  
30 amino acid fragment corresponding to amino acid residues 102-206 of the amino acid sequence shown in SEQ ID NO: 10.

In another embodiment the invention relates to a variant of a parent Termamyl-like  $\alpha$ -amylase, in which variant at least one of the amino acid residues of the parent  $\alpha$ -amylase, which is/are present in a fragment corresponding to the amino acid fragment 117-181 of the amino acid sequence of SEQ ID NO: 4, has/have been deleted or replaced with one or more of the amino acid residues, which is/are present in an amino acid fragment corresponding to the amino acid fragment 98-206 of the amino acid sequence shown in SEQ ID NO: 10, or in which one or more additional amino acid residues has been added using the relevant part of SEQ ID NO: 10 or a corresponding part of another Fungamyl-like  $\alpha$ -amylase as a template.

For instance, the variant may be one, in which the amino acid fragment X-Y of the parent  $\alpha$ -amylase, which corresponds to or is within the amino acid fragment 117-177 of SEQ ID NO: 4, has/have been replaced with an amino acid fragment Z-V, which corresponds to or is within the amino acid fragment 98-202 of the amino acid sequence shown in SEQ ID NO: 10, in which variant

X is an amino acid residue corresponding to the amino acid occupying position 117, 118, 119, 120 or 121 of SEQ ID NO: 4,

Y is an amino acid residue corresponding to the amino acid occupying position 174, 175, 176 or 177 of SEQ ID NO: 4,

Z is an amino acid residue corresponding to the amino acid occupying position 98, 99, 100, 101, 102 of SEQ ID NO: 10, and

V is an amino acid residue corresponding to the amino acid occupying position 199, 200, 201 or 202 of SEQ ID NO: 10.

A specific example of a variant according to this embodiment of the invention is a variant, in which the amino acid fragment of the parent  $\alpha$ -amylase, which corresponds to amino acid residues 121-174 of SEQ ID NO: 4, has been replaced with the amino acid fragment corresponding to amino acid residues 102-199 of the amino acid sequence shown in SEQ ID NO: 10.

#### *Loop 1 modifications - minimal addition*

In a further embodiment the present invention relates to a variant of a parent Termamyl-like  $\alpha$ -amylase, in which variant at least one of the amino acid residues of the parent  $\alpha$ -amylase, which is/are present in an amino acid fragment corresponding to the amino



acid fragment 12-19 of the amino acid sequence of SEQ ID NO: 4, has/have been deleted or replaced with one or more of the amino acid residues, which is/are present in an amino acid fragment which corresponds to the amino acid fragment 28-42 of SEQ ID NO: 10, or in which one or more additional amino acid residues has/have been inserted using the relevant part of SEQ ID NO: 10 or a corresponding part of another Fungamyl-like  $\alpha$ -amylase as a template.

For instance, the variant may be one, in which the amino acid fragment X-Y of the parent  $\alpha$ -amylase, which corresponds to or is within the amino acid fragment 12-19 of SEQ ID NO: 4, has/have been replaced with an amino acid fragment Z-V, which corresponds to or is within the amino acid fragment 28-42 of the amino acid sequence shown in SEQ ID NO: 10, in which variant

X is an amino acid residue corresponding to the amino acid occupying position 12, 13 or 14 of SEQ ID NO: 4,

Y is an amino acid residue corresponding to the amino acid occupying position 15, 16, 17, 18 or 19 of SEQ ID NO: 4,

Z is an amino acid residue corresponding to the amino acid occupying position 28, 29, 30, 31 or 32 of SEQ ID NO: 10, and

V is an amino acid residue corresponding to the amino acid occupying position 38, 39, 40, 41 or 42 of SEQ ID NO: 10.

A specific example of a variant according to this aspect of the invention is a variant, in which the amino acid fragment of the parent  $\alpha$ -amylase, which corresponds to amino acid residues 14-15 of SEQ ID NO: 4, has been replaced with the amino acid fragment corresponding to amino acid residues 32-38 of the amino acid sequence shown in SEQ ID NO: 10.

#### *Loop 1 modifications - complete loop*

In a further embodiment the invention relates to a variant of a parent Termamyl-like  $\alpha$ -amylase, in which variant at least one of the amino acid residues of the parent  $\alpha$ -amylase, which is present in a fragment corresponding to amino acid residues 7-23 of the amino acid sequence of SEQ ID NO: 4, has/have been deleted or replaced with one or more amino acid residues, which is/are present in an amino acid fragment corresponding to amino acid residues 13-45 of the amino acid sequence shown in SEQ ID NO: 10, or in which

one or more additional amino acid residues has/have been inserted using the relevant part of SEQ ID NO: 10 or a corresponding part of another Fungamyl-like  $\alpha$ -amylase as a template.

For instance, the variant may be one, in which the amino acid fragment X-Y of the parent  $\alpha$ -amylase, which corresponds to or is within the amino acid fragment 7-23 of SEQ ID NO: 4, has/have been replaced with an amino acid fragment Z-V, which corresponds to or is within the amino acid fragment 13-45 of the amino acid sequence shown in SEQ ID NO: 10, in which variant

X is an amino acid residue corresponding to the amino acid occupying position 7 or 8 of SEQ ID NO: 4,

Y is an amino acid residue corresponding to the amino acid occupying position 18, 19, 20, 21, 22 or 23 of SEQ ID NO: 4,

Z is an amino acid residue corresponding to the amino acid occupying position 13 or 14 of SEQ ID NO: 10, and

V is an amino acid residue corresponding to the amino acid occupying position 40, 41, 42, 43, 44 or 45 of SEQ ID NO: 10.

A specific variant according to this embodiment is one, in which the amino acid fragment of the parent  $\alpha$ -amylase, which corresponds to amino acid residues 8-18 of SEQ ID NO: 4, has been replaced with the amino acid fragment corresponding to amino acid residues 14-40 of the amino acid sequence shown in SEQ ID NO: 10.

#### *Loop 8 modifications*

In a further embodiment the invention relates to a variant of a parent Termamyl-like  $\alpha$ -amylase, in which variant at least one of the amino acid residues of the parent  $\alpha$ -amylase, which is present in a fragment corresponding to amino acid residues 322-346 of the amino acid sequence of SEQ ID NO: 2, has/have been deleted or replaced with one or more amino acid residues, which is/are present in an amino acid fragment corresponding to amino acid residues 291-313 of the amino acid sequence shown in SEQ ID NO: 10, or in which one or more additional amino acid residues has/have been inserted using the relevant part of SEQ ID NO: 10 or a corresponding part of another Fungamyl-like  $\alpha$ -amylase as a template.

For instance, the variant may be one, in which the amino acid fragment X-Y of the parent  $\alpha$ -amylase, which corresponds to or is within the amino acid fragment 322-346

of SEQ ID NO: 2, has/have been replaced with an amino acid fragment Z-V, which corresponds to or is within the amino acid fragment 291-313 of the amino acid sequence shown in SEQ ID NO: 10, in which variant

X is an amino acid residue corresponding to the amino acid occupying position 322, 323, 324 or 325 of SEQ ID NO: 2,

Y is an amino acid residue corresponding to the amino acid occupying position 343, 344, 345 or 346 of SEQ ID NO: 2,

Z is an amino acid residue corresponding to the amino acid occupying position 291, 292, 293 or 294 of SEQ ID NO: 10, and

V is an amino acid residue corresponding to the amino acid occupying position 310, 311, 312 or 313 of SEQ ID NO: 10.

A specific variant according to this aspect of the invention is one, in which the amino acid fragment of the parent  $\alpha$ -amylase, which corresponds to amino acid residues 325-345 of SEQ D No. 2, has been replaced with the amino acid fragment corresponding to amino acid residues 294-313 of the amino acid sequence shown in SEQ ID NO: 10.

#### Ca<sup>2+</sup> dependency

It is highly desirable to be able to decrease the Ca<sup>2+</sup> dependency of a Termamyl-like  $\alpha$ -amylase. Accordingly, in a further aspect the invention relates to a variant of a parent Termamyl-like  $\alpha$ -amylase, which exhibits  $\alpha$ -amylase activity and which has a decreased Ca<sup>2+</sup> dependency as compared to the parent  $\alpha$ -amylase. The decreased Ca<sup>2+</sup> dependency has the functional result that the variant exhibits a satisfactory amylolytic activity in the presence of a lower concentration of calcium ion in the extraneous medium than is necessary for the parent enzyme and, for example, therefore is less sensitive than the parent to calcium ion-depleting conditions such as those obtained in media containing calcium-complexing agents (such as certain detergent builders).

The decreased Ca<sup>2+</sup> dependency of the variant of the invention may advantageously be achieved by increasing the Ca<sup>2+</sup> binding affinity of the parent Termamyl-like  $\alpha$ -amylase, in other words the stronger the Ca<sup>2+</sup> binding of the enzyme, the lower is the Ca<sup>2+</sup> dependency.

It is presently believed that amino acid residues located within 10Å from a sodium or calcium ion are involved in or are of importance for the Ca<sup>2+</sup> binding capability of the enzyme.

Accordingly, the variant according to this aspect of the invention is preferably one, which has been modified in one or more amino acid residues present within 10Å from a calcium and/or sodium ion identified in the three-dimensional Termamyl-like α-amylase structure in such a manner that the affinity of the α-amylase for calcium is increased.

The amino acid residues found within a distance of 10Å from the Ca<sup>2+</sup> binding sites of the *B. licheniformis* α-amylase with the amino acid sequence SEQ ID NO: 2 were determined as described in Example 2 and are as follows:

V102, I103, N104, H105, K106, R125, W155, W157, Y158, H159, F160, D161, G162, T163, Y175, K176, F177, G178, K180, A181, W182, D183, W184, E185, V186, S187, N192, Y193, D194, Y195, L196, M197, Y198, A199, D200, I201, D202, Y203, D204, H205, P206, V208, A209, D231, A232, V233, K234, H235, I236, K237, F238, F240, L241, A294, A295, S296, T297, Q298, G299, G300, G301, Y302, D303, M304, R305, K306, L307, W342, F343, L346, Q393, Y394, Y396, H405, H406, D407, I408, V409, R413, E414, G415, D416, S417, V419, A420, N421, S422, G423, L424, I428, T429, D430, G431, P432, V440, G441, R442, Q443, N444, A445, G446, E447, T448, W449, I462, G475, Y480, V481, Q482, R483.

In order to construct a variant according to this aspect of the invention it is desirable to replace at least one of the above mentioned amino acid residues (or an amino acid residue occupying an equivalent position in another Termamyl-like α-amylase than that defined by SEQ ID NO: 2), which is contemplated to be involved in providing a non-optimal calcium binding, with any other amino acid residue which improves the Ca<sup>2+</sup> binding affinity of the variant enzyme. In practice, the identification and subsequent modification of the amino acid residue is performed by the following method:

i) identifying an amino acid residue within 10Å from a Ca<sup>2+</sup> binding site of a Termamyl-like α-amylase structure, which from structural or functional considerations is believed to be responsible for a non-optimal calcium ion interaction,

ii) constructing a variant in which said amino acid residue is replaced with another amino acid residue which from structural or functional considerations is believed to be important for establishing a higher Ca<sup>2+</sup> binding affinity, and testing the Ca<sup>2+</sup> dependency of the resulting Termamyl-like α-amylase variant.

In the present context, the term "non-optimal calcium ion interaction" is intended to indicate that the amino acid residue in question is selected on the basis of a presumption that substituting said amino acid residue for another may improve a calcium ion binding interaction of the enzyme. For instance, the amino acid residue in question may be selected on the basis of one or more of the following considerations:

- to obtain an improved interaction between a calcium ion and an amino acid residue located near to the surface of the enzyme (as identified from the structure of the Termamyl-like  $\alpha$ -amylase). For instance, if the amino acid residue in question is exposed to a surrounding solvent, it may be advantageous to increase the shielding of said amino acid residue from the solvent so as to provide for a stronger interaction between said amino acid residue and a calcium ion. This can be achieved by replacing said residue (or an amino acid residue in the vicinity of said residue contributing to the shielding) by an amino acid residue which is more bulky or otherwise results in an improved shielding effect.

- to stabilize a calcium binding site, for instance by stabilizing the structure of the Termamyl-like  $\alpha$ -amylase (e.g. by stabilizing the contacts between the A, B and C domains or stabilizing one or more of the domains as such). This may, e.g., be achieved by providing for a better coordination to amino acid side chains, which may, e.g., be obtained by replacing an N residue with a D residue and/or a Q residue with an E residue (e.g. N104D), e.g. within 10Å, and preferably within 3 or 4Å, of a calcium binding site.

- to protect the calcium binding site or to improve the coordination between the calcium ion and the calcium binding site, e.g. by providing a stronger interaction between the ion and the binding site.

Before actually constructing a Termamyl-like  $\alpha$ -amylase variant according to the above principles it may be convenient to evaluate the contemplated amino acid modification by its accommodation into the Termamyl-like  $\alpha$ -amylase structure, e.g. into a model structure of the parent Termamyl-like  $\alpha$ -amylase.

Preferably, the amino acid residue to be modified is located within 8Å of a  $\text{Ca}^{2+}$  binding site residue, such as within 5Å of such residue. The amino acid residues within 8Å and 5Å, respectively, may easily be identified by an analogous method used for identifying amino acid residues within 10Å (cf. Example 2).

The following mutation is contemplated to be of particular interest with respect to decreasing the  $\text{Ca}^{2+}$  dependency of a Termamyl-like  $\alpha$ -amylase:

N104D (of the *B. licheniformis*  $\alpha$ -amylase SEQ ID NO: 2, or an equivalent (N to D) mutation of an equivalent position in another Termamyl-like  $\alpha$ -amylase.)

In connection with substitutions of relevance for  $\text{Ca}^{2+}$  dependency, some other substitutions appear to be of importance in stabilizing the enzyme conformation (for instance the Domains A-B and/or Domains A-C interactions contributing to the overall stability of the enzyme) in that they may, e.g., enhance the strength of binding or retention of calcium ion or sodium ion at or within a calcium or sodium binding site, respectively, within the parent Termamyl-like  $\alpha$ -amylase.

It is desirable to stabilize the C-domain in order to increase the calcium stability and/or thermostability of the enzyme. In this connection the stabilization may result in a stabilization of the binding of calcium by the enzyme, and an improved contact between the C-domain and the A-domain (of importance for thermostability). The latter may be achieved by introduction of cystein bridges, salt bridges or increase hydrogen, hydrophobic and/or electrostatic interactions.

For instance, the C-domain of the *B. licheniformis*  $\alpha$ -amylase having the amino acid sequence shown in SEQ ID NO: 2 may be stabilized by introduction of a cystein bridge between domain A and domain C, e.g. by introducing of the following mutations: A349C+I479C and/or L346C+I430C.

A salt bridge may be obtained by introduction of the following mutations:

N457D,E

N457D,E+K385R

F350D,E+I430R,K

F350D,E+I411R,K

The calcium site of Domain C may be stabilized by replacing the amino acid residues H408 and/or G303 with any other amino acid residue. Of particular interest is the following mutations:

H408Q,E,N,D and/or G303N,D,Q,E

which are contemplated to provide a better calcium binding or protection from calcium depletion.

Similar mutations may be introduced in equivalent positions of other Termamyl-like  $\alpha$ -amylases.

Other substitution mutations (relative to *B. licheniformis*  $\alpha$ -amylase, SEQ ID NO: 2) which appear to be of importance, *inter alia*, in the context of reducing calcium dependency include the following:

In Domain A:

R23K;  
A209V; and  
G310D;

In Domain B:

H156Y;

A181E,D,Q,N,V,T (which appear to result in shielding of the outermost  $\text{Ca}^{2+}$  binding site in the junction region between Domain A and Domain B to some extent);

I201(bulkier amino acid), e.g. I201W,F,L (which appear to result in slight alterations in the geometry of the region in the immediate vicinity of the  $\text{Ca}^{2+}$ - $\text{Na}^{+}$ - $\text{Ca}^{2+}$  binding site(s) in the junction region between Domain A and Domain B, and in the geometry and/or size of a nearby hole/cavity); and

Y203E,Q (which are believed to result in stronger binding of the outermost  $\text{Ca}^{2+}$  ion in its binding site in the junction region between Domain A and Domain B);  
(or equivalent mutations in equivalent positions in another Termamyl-like  $\alpha$ -amylase).

Substitutions of R214 and P345 of *B. licheniformis*  $\alpha$ -amylase, SEQ ID NO: 2 (or equivalent mutations in equivalent positions in another Termamyl-like  $\alpha$ -amylase) with other amino acids may also be of importance in this connection.

Variants with altered activity at higher/lower pH

It is contemplated that it is possible to change the pH optima of a Termamyl-like  $\alpha$ -amylase or the enzymatic activity at a given pH by changing the pKa of the active site residues. This may be achieved, e.g. by changing the electrostatic interaction or hydrophobic interaction between functional groups of amino acid side chains of the amino acid residue to be modified and of its close surroundings. This may, e.g., be accomplished by the following method:

i) in a structure of the Termamyl-like  $\alpha$ -amylase in question to identifying an amino acid residue within 15Å from an active site residue, in particular 10Å from an active

site residue, which amino acid residue is contemplated to be involved in electrostatic or hydrophobic interactions with an active site residue,

ii) replacing, in the structure, said amino acid residue with an amino acid residue which changes the electrostatic and/or hydrophobic surroundings of an active site residue and evaluating the accommodation of the amino acid residue in the structure,

iii) optionally repeating step i) and/or ii) until an amino acid replacement has been identified which is accommodated into the structure,

iv) constructing a Termamyl-like  $\alpha$ -amylase variant resulting from steps i), ii) and optionally iii) and testing the pH dependent enzymatic activity of interest of said variant.

In the above method it may be of particular relevance to add a positively charged residue within 5 Å of a glutamate (thereby lowering the pKa of the glutamate from about 4.5 to 4), or to add a negatively charged residue within 5 Å of a glutamate (thereby increasing the pKa to about 5), or to make similar modifications within a distance of about 5 Å of a Histidine.

On the basis of electrostatic considerations [see, e.g., M.K. Gilson, Current Opinion in Structural Biology 5 (1995) pp. 216-223; and B. Honig and A. Nicholls, Science 268 (1995) pp. 1144-1149; and references given therein] and hygroscopicity considerations in relation to the three dimensional structure of a Termamyl-like  $\alpha$ -amylase disclosed herein, mutations of relevance, *inter alia*, for changing (increasing or decreasing) the pH optimum of a Termamyl-like  $\alpha$ -amylase are believed to include the following mutations or equivalents thereof [referring here to the sequence of *B. licheniformis*  $\alpha$ -amylase (SEQ ID NO: 2)]:

Q9K,L,E; F11R,K,E; E12Q; D100N,L; V101H,R,K,D,E,F; V102A,T; I103H,K; N104R,K,D; H105R,K,D,E,W,F; L196R,K,D,E,F,Y; I212R,K,D,E; L230H,K,I; A232G,H,F,S,V; V233D; K234L,E; I236R,K,N,H,D,E; L241R,K,D,E,F; A260S; W263H; Q264R,D,K,E; N265K,R,D; A269R,K,D,E; L270R,K,H,D,E; V283H,D; F284H; D285N,L; V286R,K,H,D,E; Y290R,E; V312R,K,D,E; F323H; D325N; N326K,H,D,L; H327Q,N,E,D,F; Q330L,E; G332D; Q333R,K,H,E,L; S334A,V,T,L,I,D; L335G,A,S,T,N; E336R+R375E; T337D,K; T338D,E; T339D; Q360K,R,E; D365N; G371D,R.

In a further aspect the invention relates to a variant of a Termamyl-like  $\alpha$ -amylase which exhibits a higher activity at a lower pH (e.g. compared to the pH optimum) than the parent  $\alpha$ -amylase. In particular, the variant comprises a mutation of an amino acid



residue corresponding to at least one of the following positions of the *B. licheniformis*  $\alpha$ -amylase (SEQ ID NO: 2):

E336, Q333, P331, I236, V102, A232, I103, L196

The following mutations are of particular interest:

5 E336R,K

Q333R,K

P331R,K

V102R,K,A,T,S,G;

I236K,R,N;

10 I103K,R;

L196K,R;

A232T,S,G;

or any combination of two or more of these variants or any combination of one or more of these variants with any of the other variants disclosed herein.

15 In a still further aspect the invention relates to a variant of a Termamyl-like  $\alpha$ -amylase which has a higher activity at a higher pH than the parent  $\alpha$ -amylase. In particular, the variant comprises a mutation of an amino acid residue corresponding to at least one of the following positions of the *B. licheniformis*  $\alpha$ -amylase (SEQ ID NO: 2):

N236, H281, Y273

20 In particular, the variant comprises a mutation corresponding to at least one of the following mutations of the *B. licheniformis*  $\alpha$ -amylase (SEQ ID NO: 2):

N326I,Y,F,L,V

H281F,I,L

Y273F,W

25 or any combination of two or more of these variants or any combination of one or more of these variants with any of the other variants disclosed herein.

A mutation which appears to be importance in relation to the specific activity of variants of the invention is a mutation corresponding to the substitution S187D in *B. licheniformis*  $\alpha$ -amylase (SEQ ID NO: 2).

### Variants with increased thermostability and/or altered temperature optimum

In a further desired aspect the invention relates to a variant of a parent Termamyl-like  $\alpha$ -amylase, which variant is the result of one or more amino acid residues having been deleted from, replaced or added to the parent  $\alpha$ -amylase so as to obtain an increased thermostability of the variant.

The Termamyl-like  $\alpha$ -amylase structure contains a number of unique internal holes, which may contain water, and a number of crevices. In order to increase the thermostability of the  $\alpha$ -amylase it may be desirable to reduce the number of holes and crevices (or reduce the size of the holes or crevices), e.g. by introducing one or more hydrophobic contacts, preferably achieved by introducing bulkier residues, in the vicinity or surroundings of the hole. For instance, the amino acid residues to be modified are those which are involved in the formation of the hole.

Accordingly, in a further aspect the present invention relates to a method of increasing the thermostability and/or altering the temperature optimum of a parent Termamyl-like  $\alpha$ -amylase, which method comprises

- i) identifying an internal hole or a crevice of the parent Termamyl-like  $\alpha$ -amylase in the three-dimensional structure of said  $\alpha$ -amylase,
- ii) replacing, in the structure, one or more amino acid residues in the neighbourhood of the hole or crevice identified in i) with another amino acid residue which from structural or functional considerations is believed to increase the hydrophobic interaction and to fill out or reduce the size of the hole or crevice,
- iii) constructing a Termamyl-like  $\alpha$ -amylase variant resulting from step ii) and testing the thermostability and/or temperature optimum of the variant.

The structure used for identifying the hole or crevice of the parent Termamyl-like  $\alpha$ -amylase may be the structure identified in Appendix 1 or a model structure of the parent Termamyl-like  $\alpha$ -amylase built thereon.

It will be understood that the hole or crevice is identified by the amino acid residues surrounding the hole/crevice, and that modification of said amino acid residues are of importance for filling or reducing the size of the hole/crevice. The particular amino acid residues referred to below are those which in crystal structure have been found to flank the hole/crevice in question.

In order to fill (completely or partly) a major hole located between domain A and B, mutation to any other amino acid residue of an amino acid residue corresponding to one or more of the following residues of the *B. licheniformis*  $\alpha$ -amylase (SEQ ID NO: 2) is contemplated:

5 L61, Y62, F67, K106, G145, I212, S151, R214, Y150, F143, R146

Of particular interest is a mutation to a more bulky amino acid residue than the amino acid residue of the parent enzyme.

Of particular interest is a variant of a Termamyl-like  $\alpha$ -amylase which comprises a mutation corresponding to the following mutations (using the numbering of *B. licheniformis*  $\alpha$ -amylase (SEQ ID NO: 2):

10 L61W,V,F;

Y62W;

F67W;

K106R,F,W;

15 G145F,W

I212F,L,W,Y,R,K;

S151 replaced with any other amino acid residue and in particular with F,W,I or L;

R214W;

Y150R,K;

20 F143W; and/or

R146W.

In order to fill a hole in the vicinity of the active site mutation to any other amino acid residue of an amino acid residue corresponding to one or more of the following residues of the *B. licheniformis*  $\alpha$ -amylase (SEQ ID NO: 2) is contemplated:

25 L241, I236.

Of interest is a mutation to a more bulky amino acid residue.

Of particular interest is a variant of a Termamyl-like  $\alpha$ -amylase which comprises a mutation corresponding to one or more of the following mutations in the *B. licheniformis*  $\alpha$ -amylase:

30 L241I,F,Y,W; and/or

I236L,F,W,Y

In order to fill a hole in the vicinity of the active site mutation to any other amino acid residue of an amino acid residue corresponding to one or more of the following residues of the *B. licheniformis*  $\alpha$ -amylase (SEQ ID NO: 2) is contemplated:

L7, V259, F284

5 Of interest is a mutation to a more bulky amino acid residue.

Of particular interest is a variant of a Termamyl-like  $\alpha$ -amylase which comprises a mutation corresponding to one or more of the following mutations in the *B. licheniformis*  $\alpha$ -amylase:

L7F,I,W

10 V259F,I,L

F284W

In order to fill a hole in the vicinity of the active site mutation to any other amino acid residue of an amino acid residue corresponding to one or more of the following residues of the *B. licheniformis*  $\alpha$ -amylase (SEQ ID NO: 2) is contemplated:

15 F350, F343

Of interest is a mutation to a more bulky amino acid residue.

Of particular interest is a variant of a Termamyl-like  $\alpha$ -amylase which comprises a mutation corresponding to one or more of the following mutations in the *B. licheniformis*  $\alpha$ -amylase:

20 F350W

F343W

In order to fill a hole in the vicinity of the active site mutation to any other amino acid residue of an amino acid residue corresponding to one or more of the following residues of the *B. licheniformis*  $\alpha$ -amylase (SEQ ID NO: 2) is contemplated:

25 L427, V481

Of interest is a mutation to a more bulky amino acid residue.

Of particular interest is a variant of a Termamyl-like  $\alpha$ -amylase which comprises a mutation corresponding to one or more of the following mutations in the *B. licheniformis*  $\alpha$ -amylase:

30 L427F,L,W

V481,F,I,L,W

It can be seen from an alignment of the amino acid sequences of  $\alpha$ -amylases from various *Bacillus* species that *B. licheniformis*  $\alpha$ -amylase and *B. amyloliquefaciens*  $\alpha$ -amylase both contain an "insertion" of three amino acids relative to, e.g., *B. stearothermophilus*  $\alpha$ -amylase.

From a model of the structure of *B. licheniformis*  $\alpha$ -amylase built on the basis of the three-dimensional structure of the Termamyl-like  $\alpha$ -amylase disclosed herein (*vide supra*), taking into account the homology of *B. licheniformis*  $\alpha$ -amylase to the Termamyl-like  $\alpha$ -amylase in question, it can be seen that the above-mentioned "insertion" lies within loop 8 (*vide supra*), making this loop bulkier in *B. licheniformis*  $\alpha$ -amylase than in the Termamyl-like  $\alpha$ -amylase and resulting in a loop that protrudes from the structure, thereby possibly destabilizing the structure. It is therefore contemplated that deletion of one or more amino acids in the region in question in *B. licheniformis* or *B. amyloliquefaciens*  $\alpha$ -amylase will improve the thermostability of these  $\alpha$ -amylases.

Especially interesting in this connection is deletion of three amino acids within the partial sequence from T369 to I377 (referring to the sequence of *B. licheniformis*  $\alpha$ -amylase), i.e. the partial sequence: T369-K370-G371-D372-S373-Q374-R375-E376-I377 (or the corresponding partial sequence in *B. amyloliquefaciens*  $\alpha$ -amylase). In addition to such deletions, substitution of one or more of the undeleted amino acids within the latter partial sequence may also be advantageous.

Preferable deletions of three amino acids in the partial sequence from T369 to I377 (in *B. licheniformis*  $\alpha$ -amylase) are deletion of K370+G371+D372 (i.e. K370\*+G371\*+D372\*) or deletion of D372+S373+Q374 (i.e. D372\*+S373\*+Q374\*) (or equivalent deletions in the corresponding partial sequence in *B. amyloliquefaciens*  $\alpha$ -amylase).

Another type of mutation which would appear to be of value in improving the thermostability of these  $\alpha$ -amylases is substitution (replacement) of the entire partial amino acid sequence from T369 to I377 (referring to the sequence of *B. licheniformis*  $\alpha$ -amylase) with one of the following partial sequences of six amino acids (sequence numbering increasing from left to right): I-P-T-H-S-V; I-P-T-H-G-V; and I-P-Q-Y-N-I (or one of the same substitutions of the corresponding partial sequence in *B. amyloliquefaciens*  $\alpha$ -amylase).

### Variants with an altered cleavage pattern

In the starch liquefaction process it is desirable to use an  $\alpha$ -amylase which is capable of degrading the starch molecules into long branched oligo saccharides (like, e.g. the Fungamyl-like  $\alpha$ -amylases) rather than shorter branched oligo saccharides (like conventional Termamyl-like  $\alpha$ -amylases). The resulting very small branched oligosaccharides (panose precursors) cannot be hydrolyzed properly by pullulanases, which in the liquefaction process are used after the  $\alpha$ -amylases and before the amyloglucosidases. Thus, in the presence of panose precursors the action of amylo-glucoamylase ends up with a high degree of the small branched limiting-dextrin, the trisaccharide panose. The presence of panose lowers the saccharification yield significantly and is thus undesirable.

Thus, one aim of the present invention is to change the degradation characteristics of a Termamyl-like  $\alpha$ -amylase to that of a Fungamyl-like  $\alpha$ -amylases without at the same time reducing the thermostability of the Termamyl-like  $\alpha$ -amylase.

Accordingly, in a further aspect the invention relates to a variant of a Termamyl-like  $\alpha$ -amylase which has a reduced ability to cleave a substrate close to the branching point.

The variant may suitably be constructed by a method which comprises

i) identifying the substrate binding area of the parent Termamyl-like  $\alpha$ -amylase in a model of the three-dimensional structure of said  $\alpha$ -amylase, (e.g. within a sphere of 4Å from the substrate binding site (as defined in the section above entitled "Substrate Binding Site"),

ii) replacing, in the model, one or more amino acid residues of the substrate binding area of the cleft identified in i), which is/are believed to be responsible for the cleavage pattern of the parent  $\alpha$ -amylase, with another amino acid residue which from structural considerations is believed to result in an altered substrate cleavage pattern, or deleting one or more amino acid residues of the substrate binding area contemplated to introduce favourable interactions to the substrate or adding one or more amino acid residues to the substrate binding area contemplated to introduce favourable interactions to the substrate, and

iii) constructing a Termamyl-like  $\alpha$ -amylase variant resulting from step ii) and testing the substrate cleavage pattern of the variant.

Of particular interest is a variant which cleaves an amylopectin substrate, from the reducing end, more than one glucose unit from the branching point, preferably more than two or three glucose units from the branching point, i.e. at a further distance from the branching point than that obtained by use of a wild type *B. licheniformis*  $\alpha$ -amylase.

Residues of particular interest in connection with this aspect of the invention correspond to the following residues of the *B. licheniformis*  $\alpha$ -amylase (SEQ ID NO: 2): V54, D53, Y56, Q333, G57, and the variants according to this aspect preferably comprises a mutation in one or more of these residues.

In particular, the variant comprises at least one of the following mutations, which are expected to prevent cleavage close to the branching point:

V54L,I,F,Y,W,R,K,H,E,Q

D53L,I,F,Y,W

Y56W

Q333W

G57all possible amino acid residues

A52amino acid residues larger than A, e.g. A52W,Y,L,F,I.

#### Variants of a fungal $\alpha$ -amylase

In a still further embodiment the invention relates to a variant of a parent Fungamyl-like  $\alpha$ -amylase, in which variant at least one of the amino acid residues of the parent  $\alpha$ -amylase, which is/are present in an amino acid fragment corresponding to amino acid residues 291-313 of the amino acid sequence of SEQ ID NO: 10, has/have been deleted or replaced with one or more of the amino acid residues, which is/are present in an amino acid fragment corresponding to amino acid residues 98-210 of the amino acid sequence shown in SEQ ID NO: 4, or in which one or more additional amino acid residues has/have been inserted using the relevant part of SEQ ID NO: 4 or a corresponding part of another Termamyl-like  $\alpha$ -amylase as a template.

For instance, the variant may be one, in which the amino acid fragment X-Y of the parent  $\alpha$ -amylase, which corresponds to or is within the amino acid fragment 117-185 of SEQ ID NO: 10, has/have been replaced with an amino acid fragment Z-V, which corresponds to or is within the amino acid fragment 98-210 of the amino acid sequence shown in SEQ ID NO: 4, in which variant

X is an amino acid residue corresponding to the amino acid occupying position 117, 118, 119, 120 or 121 of SEQ ID NO: 10,

Y is an amino acid residue corresponding to the amino acid occupying position 181, 182, 183, 184 or 185 of SEQ ID NO: 10,

5 Z is an amino acid residue corresponding to the amino acid occupying position 98, 99, 100, 101 or 102 of SEQ ID NO: 4, and

V is an amino acid residue corresponding to the amino acid occupying position 206, 207, 208, 209 or 210 of SEQ ID NO: 4.

10 A specific example of a variant according to this aspect of the invention is one, in which the amino acid fragment of the parent  $\alpha$ -amylase, which corresponds to amino acid residues 121-181 of SEQ ID NO: 10, has been replaced with the amino acid fragment corresponding to amino acid residues 102-206 of the amino acid sequence shown in SEQ ID NO: 4.

15 Another example of a variant according to this aspect of the invention is one, in which the amino acid fragment of the parent  $\alpha$ -amylase, which corresponds to amino acid residues 121-174 of SEQ ID NO: 10, has been replaced with the amino acid fragment corresponding to amino acid residues 102-199 of the amino acid sequence shown in SEQ ID NO: 4.

20 In a further embodiment the invention relates to a variant of a parent Fungamyl-like  $\alpha$ -amylase, in which an amino acid fragment corresponding to amino acid residues 181-184 of the amino acid sequence shown in SEQ ID NO: 10 has been deleted.

#### General mutations in variants of the invention

25 It may be preferred that the variant of the invention or prepared in accordance with the method of the invention comprises one or more modifications in addition to those outlined above. Thus, it may be advantageous that one or more proline residues present in the part of the  $\alpha$ -amylase variant having been modified is/are replaced with a non-proline residue which may be any of the possible, naturally occurring non-proline residues, and which preferably is an alanine, glycine, serine, threonine, valine or leucine.

30 Analogously, it may be preferred that one or more cysteine residues present in the amino acid residues with which the parent  $\alpha$ -amylase is modified are replaced with a non-cysteine residues such as serine, alanine, threonine, glycine, valine or leucine.



Furthermore, the variant of the invention may, as the only modification or in combination with any of the above outlined modifications, be modified so that one or more Asp and/or Glu present in an amino acid fragment corresponding to the amino acid sequence fragment 185-209 of SEQ ID NO: 2 is/are replaced by an Asn and/or Gln, respectively. Also of interest is the replacement of one or more of the Lys residues present in the Termamyl-like  $\alpha$ -amylase by an Arg residue.

It will be understood that in accordance with the present invention variants may be prepared which carry two or more of the above outlined modifications. For instance, variants may be prepared which comprises a modification in the loop 1 and loop 2 region, a modification in loop 2 and limited loop 3, a modification in loop 1, loop 2, loop 3 and loop 8, etc.

Furthermore, it may be advantageous to introduce point-mutations in any of the variants described herein.

#### Methods of preparing $\alpha$ -amylase variants

Several methods for introducing mutations into genes are known in the art. After a brief discussion of the cloning of  $\alpha$ -amylase-encoding DNA sequences, methods for generating mutations at specific sites within the  $\alpha$ -amylase-encoding sequence will be discussed.

#### Cloning a DNA sequence encoding an $\alpha$ -amylase

The DNA sequence encoding a parent  $\alpha$ -amylase may be isolated from any cell or microorganism producing the  $\alpha$ -amylase in question, using various methods well known in the art. First, a genomic DNA and/or cDNA library should be constructed using chromosomal DNA or messenger RNA from the organism that produces the  $\alpha$ -amylase to be studied. Then, if the amino acid sequence of the  $\alpha$ -amylase is known, homologous, labelled oligonucleotide probes may be synthesized and used to identify  $\alpha$ -amylase-encoding clones from a genomic library prepared from the organism in question. Alternatively, a labelled oligonucleotide probe containing sequences homologous to a known  $\alpha$ -amylase gene could be used as a probe to identify  $\alpha$ -amylase-encoding clones, using hybridization and washing conditions of lower stringency.

Yet another method for identifying  $\alpha$ -amylase-encoding clones would involve inserting fragments of genomic DNA into an expression vector, such as a plasmid, transforming  $\alpha$ -amylase-negative bacteria with the resulting genomic DNA library, and then plating the transformed bacteria onto agar containing a substrate for  $\alpha$ -amylase, thereby allowing clones expressing the  $\alpha$ -amylase to be identified.

Alternatively, the DNA sequence encoding the enzyme may be prepared synthetically by established standard methods, e.g. the phosphoroamidite method described by S.L. Beaucage and M.H. Caruthers (1981) or the method described by Matthes et al. (1984). In the phosphoroamidite method, oligonucleotides are synthesized, e.g. in an automatic DNA synthesizer, purified, annealed, ligated and cloned in appropriate vectors.

Finally, the DNA sequence may be of mixed genomic and synthetic origin, mixed synthetic and cDNA origin or mixed genomic and cDNA origin, prepared by ligating fragments of synthetic, genomic or cDNA origin (as appropriate, the fragments corresponding to various parts of the entire DNA sequence), in accordance with standard techniques. The DNA sequence may also be prepared by polymerase chain reaction (PCR) using specific primers, for instance as described in US 4,683,202 or R.K. Saiki et al. (1988).

#### Site-directed mutagenesis

Once an  $\alpha$ -amylase-encoding DNA sequence has been isolated, and desirable sites for mutation identified, mutations may be introduced using synthetic oligonucleotides. These oligonucleotides contain nucleotide sequences flanking the desired mutation sites; mutant nucleotides are inserted during oligonucleotide synthesis. In a specific method, a single-stranded gap of DNA, bridging the  $\alpha$ -amylase-encoding sequence, is created in a vector carrying the  $\alpha$ -amylase gene. Then the synthetic nucleotide, bearing the desired mutation, is annealed to a homologous portion of the single-stranded DNA. The remaining gap is then filled in with DNA polymerase I (Klenow fragment) and the construct is ligated using T4 ligase. A specific example of this method is described in Morinaga et al. (1984). US 4,760,025 discloses the introduction of oligonucleotides encoding multiple mutations by performing minor alterations of the cassette. However, an even greater variety of mutations can be introduced at any one time by the Morinaga method, because a multitude of oligonucleotides, of various lengths, can be introduced.

Another method of introducing mutations into  $\alpha$ -amylase-encoding DNA sequences is described in Nelson and Long (1989). It involves the 3-step generation of a PCR fragment containing the desired mutation introduced by using a chemically synthesized DNA strand as one of the primers in the PCR reactions. From the PCR-generated fragment, a DNA  
5 fragment carrying the mutation may be isolated by cleavage with restriction endonucleases and reinserted into an expression plasmid.

#### Random mutagenesis

Random mutagenesis is suitably performed either as localized or region-specific  
10 random mutagenesis in at least three parts of the gene translating to the amino acid sequence shown in question, or within the whole gene.

For region-specific random mutagenesis with a view to improving the thermal stability of a parent Termamyl-like  $\alpha$ -amylase, codon positions corresponding to the following amino acid residues of the *B. licheniformis*  $\alpha$ -amylase (SEQ ID NO: 2) may appropriately be targeted:

#### To improve the stability of the calcium site between Domain A and C

I428-A435

T297-L308

F403-V409  
20

#### To improve the stability between domain A and B:

D180-D204

H156-T163

A232-F238  
25

With a view to achieving improved binding of a substrate (i.e. improved binding of a carbohydrate species, such as amylose or amylopectin) by a Termamyl-like  $\alpha$ -amylase variant, modified (e.g. higher) substrate specificity and/or modified (e.g. higher) specificity with respect to cleavage (hydrolysis) of substrate, it appears that the following  
30 codon positions for the amino acid sequence shown in SEQ ID NO: 2 (or equivalent codon positions for another parent Termamyl-like  $\alpha$ -amylase in the context of the invention) may particularly appropriately be targeted:

13-18  
50-56  
70-76  
102-109  
163-172  
189-199  
229-235  
360-364  
327-335

10 The random mutagenesis of a DNA sequence encoding a parent  $\alpha$ -amylase to be performed in accordance with step a) of the above-described method of the invention may conveniently be performed by use of any method known in the art.

15 For instance, the random mutagenesis may be performed by use of a suitable physical or chemical mutagenizing agent, by use of a suitable oligonucleotide, or by subjecting the DNA sequence to PCR generated mutagenesis. Furthermore, the random mutagenesis may be performed by use of any combination of these mutagenizing agents.

The mutagenizing agent may, e.g., be one which induces transitions, transversions, inversions, scrambling, deletions, and/or insertions.

20 Examples of a physical or chemical mutagenizing agent suitable for the present purpose include ultraviolet (UV) irradiation, hydroxylamine, N-methyl-N'-nitro-N-nitrosoguanidine (MNNG), O-methyl hydroxylamine, nitrous acid, ethyl methane sulphonate (EMS), sodium bisulphite, formic acid, and nucleotide analogues.

25 When such agents are used, the mutagenesis is typically performed by incubating the DNA sequence encoding the parent enzyme to be mutagenized in the presence of the mutagenizing agent of choice under suitable conditions for the mutagenesis to take place, and selecting for mutated DNA having the desired properties.

30 When the mutagenesis is performed by the use of an oligonucleotide, the oligonucleotide may be doped or spiked with the three non-parent nucleotides during the synthesis of the oligonucleotide at the positions which are to be changed. The doping or spiking may be done so that codons for unwanted amino acids are avoided. The doped or spiked oligonucleotide can be incorporated into the DNA encoding the amylolytic enzyme by any published technique, using e.g. PCR, LCR or any DNA polymerase and ligase.

When PCR-generated mutagenesis is used, either a chemically treated or non-treated gene encoding a parent  $\alpha$ -amylase enzyme is subjected to PCR under conditions that increase the misincorporation of nucleotides (Deshler 1992; Leung et al., Technique, Vol.1, 1989, pp. 11-15).

5 A mutator strain of *E. coli* (Fowler et al., Molec. Gen. Genet., 133, 1974, pp. 179-191), *S. cerevisiae* or any other microbial organism may be used for the random mutagenesis of the DNA encoding the amylolytic enzyme by e.g. transforming a plasmid containing the parent enzyme into the mutator strain, growing the mutator strain with the plasmid and isolating the mutated plasmid from the mutator strain. The mutated plasmid may  
10 subsequently be transformed into the expression organism.

The DNA sequence to be mutagenized may conveniently be present in a genomic or cDNA library prepared from an organism expressing the parent amylolytic enzyme. Alternatively, the DNA sequence may be present on a suitable vector such as a plasmid or a bacteriophage, which as such may be incubated with or otherwise exposed to the mutagenizing agent. The DNA to be mutagenized may also be present in a host cell either by being integrated in the genome of said cell or by being present on a vector harboured in the cell. Finally, the DNA to be mutagenized may be in isolated form. It will be understood that the DNA sequence to be subjected to random mutagenesis is preferably a cDNA or a genomic DNA sequence.

15 In some cases it may be convenient to amplify the mutated DNA sequence prior to the expression step (b) or the screening step (c) being performed. Such amplification may be performed in accordance with methods known in the art, the presently preferred method being PCR-generated amplification using oligonucleotide primers prepared on the basis of the DNA or amino acid sequence of the parent enzyme.

25 Subsequent to the incubation with or exposure to the mutagenizing agent, the mutated DNA is expressed by culturing a suitable host cell carrying the DNA sequence under conditions allowing expression to take place. The host cell used for this purpose may be one which has been transformed with the mutated DNA sequence, optionally present on a vector, or one which was carried the DNA sequence encoding the parent enzyme during the  
30 mutagenesis treatment. Examples of suitable host cells are the following: grampositive bacteria such as *Bacillus subtilis*, *Bacillus licheniformis*, *Bacillus lentus*, *Bacillus brevis*, *Bacillus stearothermophilus*, *Bacillus alkalophilus*, *Bacillus amyloliquefaciens*, *Bacillus coagulans*,

*Bacillus circulans*, *Bacillus lautus*, *Bacillus megaterium*, *Bacillus thuringiensis*, *Streptomyces lividans* or *Streptomyces murinus*; and gramnegative bacteria such as *E.coli*.

The mutated DNA sequence may further comprise a DNA sequence encoding functions permitting expression of the mutated DNA sequence.

Localized random mutagenesis: the random mutagenesis may advantageously be localized to a part of the parent  $\alpha$ -amylase in question. This may, e.g., be advantageous when certain regions of the enzyme have been identified to be of particular importance for a given property of the enzyme, and when modified are expected to result in a variant having improved properties. Such regions may normally be identified when the tertiary structure of the parent enzyme has been elucidated and related to the function of the enzyme.

The localized random mutagenesis is conveniently performed by use of PCR-generated mutagenesis techniques as described above or any other suitable technique known in the art.

Alternatively, the DNA sequence encoding the part of the DNA sequence to be modified may be isolated, e.g. by being inserted into a suitable vector, and said part may subsequently be subjected to mutagenesis by use of any of the mutagenesis methods discussed above.

With respect to the screening step in the above-mentioned method of the invention, this may conveniently be performed by use of an assay based on the following principle:

A microorganism capable of expressing the mutated amylolytic enzyme of interest is incubated on a suitable medium and under suitable conditions for the enzyme to be secreted, the medium being provided with a double filter comprising a first protein-binding filter and on top of that a second filter exhibiting a low protein binding capability. The microorganism is located on the second filter. Subsequent to the incubation, the first filter comprising enzymes secreted from the microorganisms is separated from the second filter comprising the microorganisms. The first filter is subjected to screening for the desired enzymatic activity and the corresponding microbial colonies present on the second filter are identified.

The filter used for binding the enzymatic activity may be any protein binding filter e.g. nylon or nitrocellulose. The topfilter carrying the colonies of the expression

organism may be any filter that has no or low affinity for binding proteins e.g. cellulose acetate or Durapore™. The filter may be pretreated with any of the conditions to be used for screening or may be treated during the detection of enzymatic activity.

The enzymatic activity may be detected by a dye, fluorescence, precipitation, pH indicator, IR-absorbance or any other known technique for detection of enzymatic activity.

The detecting compound may be immobilized by any immobilizing agent e.g. agarose, agar, gelatine, polyacrylamide, starch, filter paper, cloth; or any combination of immobilizing agents.

α-Amylase activity is detected by Cibacron Red labelled amylopectin, which is immobilized on agarose. For screening for variants with increased thermal and high-pH stability, the filter with bound α-amylase variants is incubated in a buffer at pH 10.5 and 60° or 65°C for a specified time, rinsed briefly in deionized water and placed on the amylopectin-agarose matrix for activity detection. Residual activity is seen as lysis of Cibacron Red by amylopectin degradation. The conditions are chosen to be such that activity due to the α-amylase having the amino acid sequence shown in SEQ ID NO: 1 can barely be detected. Stabilized variants show, under the same conditions, increased colour intensity due to increased liberation of Cibacron Red.

For screening for variants with an activity optimum at a lower temperature and/or over a broader temperature range, the filter with bound variants is placed directly on the amylopectin-Cibacron Red substrate plate and incubated at the desired temperature (e.g. 4°C, 10°C or 30°C) for a specified time. After this time activity due to the α-amylase having the amino acid sequence shown in SEQ ID NO: 1 can barely be detected, whereas variants with optimum activity at a lower temperature will show increase amylopectin lysis. Prior to incubation onto the amylopectin matrix, incubation in all kinds of desired media - e.g. solutions containing Ca<sup>2+</sup>, detergents, EDTA or other relevant additives - can be carried out in order to screen for changed dependency or for reaction of the variants in question with such additives.

#### Testing of variants of the invention

The testing of variants of the invention may suitably be performed by determining the starch-degrading activity of the variant, for instance by growing host cells transformed with a DNA sequence encoding a variant on a starch-containing agarose plate and

identifying starch-degrading host cells. Further testing as to altered properties (including specific activity, substrate specificity, cleavage pattern, thermoactivation, pH optimum, pH dependency, temperature optimum, and any other parameter) may be performed in accordance with methods known in the art.

#### Expression of $\alpha$ -amylase variants

According to the invention, a DNA sequence encoding the variant produced by methods described above, or by any alternative methods known in the art, can be expressed, in enzyme form, using an expression vector which typically includes control sequences encoding a promoter, operator, ribosome binding site, translation initiation signal, and, optionally, a repressor gene or various activator genes.

The recombinant expression vector carrying the DNA sequence encoding an  $\alpha$ -amylase variant of the invention may be any vector which may conveniently be subjected to recombinant DNA procedures, and the choice of vector will often depend on the host cell into which it is to be introduced. Thus, the vector may be an autonomously replicating vector, i.e. a vector which exists as an extrachromosomal entity, the replication of which is independent of chromosomal replication, e.g. a plasmid, a bacteriophage or an extrachromosomal element, minichromosome or an artificial chromosome. Alternatively, the vector may be one which, when introduced into a host cell, is integrated into the host cell genome and replicated together with the chromosome(s) into which it has been integrated.

In the vector, the DNA sequence should be operably connected to a suitable promoter sequence. The promoter may be any DNA sequence which shows transcriptional activity in the host cell of choice and may be derived from genes encoding proteins either homologous or heterologous to the host cell. Examples of suitable promoters for directing the transcription of the DNA sequence encoding an  $\alpha$ -amylase variant of the invention, especially in a bacterial host, are the promoter of the *lac* operon of *E.coli*, the *Streptomyces coelicolor* agarase gene *dagA* promoters, the promoters of the *Bacillus licheniformis*  $\alpha$ -amylase gene (*amyL*), the promoters of the *Bacillus stearothermophilus* maltogenic amylase gene (*amyM*), the promoters of the *Bacillus amyloliquefaciens*  $\alpha$ -amylase (*amyQ*), the promoters of the *Bacillus subtilis* xylA and xylB genes etc. For transcription in a fungal host, examples of useful promoters are those derived from the gene encoding *A. oryzae* TAKA amylase, *Rhizomucor miehei* aspartic proteinase, *A. niger* neutral  $\alpha$ -amylase, *A. niger* acid stable  $\alpha$ -



amylase, *A. niger* glucoamylase, *Rhizomucor miehei* lipase, *A. oryzae* alkaline protease, *A. oryzae* triose phosphate isomerase or *A. nidulans* acetamidase.

The expression vector of the invention may also comprise a suitable transcription terminator and, in eukaryotes, polyadenylation sequences operably connected to the DNA sequence encoding the  $\alpha$ -amylase variant of the invention. Termination and polyadenylation sequences may suitably be derived from the same sources as the promoter.

The vector may further comprise a DNA sequence enabling the vector to replicate in the host cell in question. Examples of such sequences are the origins of replication of plasmids pUC19, pACYC177, pUB110, pE194, pAMB1 and pIJ702.

The vector may also comprise a selectable marker, e.g. a gene the product of which complements a defect in the host cell, such as the *dal* genes from *B. subtilis* or *B. licheniformis*, or one which confers antibiotic resistance such as ampicillin, kanamycin, chloramphenicol or tetracyclin resistance. Furthermore, the vector may comprise *Aspergillus* selection markers such as *amdS*, *argB*, *niaD* and *sC*, a marker giving rise to hygromycin resistance, or the selection may be accomplished by co-transformation, e.g. as described in WO 91/17243.

While intracellular expression may be advantageous in some respects, e.g. when using certain bacteria as host cells, it is generally preferred that the expression is extracellular. In general, the *Bacillus*  $\alpha$ -amylases mentioned herein comprise a preregion permitting secretion of the expressed protease into the culture medium. If desirable, this preregion may be replaced by a different preregion or signal sequence, conveniently accomplished by substitution of the DNA sequences encoding the respective preregions.

The procedures used to ligate the DNA construct of the invention encoding an  $\alpha$ -amylase variant, the promoter, terminator and other elements, respectively, and to insert them into suitable vectors containing the information necessary for replication, are well known to persons skilled in the art (cf., for instance, Sambrook et al. (1989)).

The cell of the invention, either comprising a DNA construct or an expression vector of the invention as defined above, is advantageously used as a host cell in the recombinant production of an  $\alpha$ -amylase variant of the invention. The cell may be transformed with the DNA construct of the invention encoding the variant, conveniently by integrating the DNA construct (in one or more copies) in the host chromosome. This integration is generally considered to be an advantage as the DNA sequence is more likely to be stably maintained

in the cell. Integration of the DNA constructs into the host chromosome may be performed according to conventional methods, e.g. by homologous or heterologous recombination. Alternatively, the cell may be transformed with an expression vector as described above in connection with the different types of host cells.

5           The cell of the invention may be a cell of a higher organism such as a mammal or an insect, but is preferably a microbial cell, e.g. a bacterial or a fungal (including yeast) cell.

10           Examples of suitable bacteria are grampositive bacteria such as *Bacillus subtilis*, *Bacillus licheniformis*, *Bacillus lentus*, *Bacillus brevis*, *Bacillus stearothermophilus*, *Bacillus alkalophilus*, *Bacillus amyloliquefaciens*, *Bacillus coagulans*, *Bacillus circulans*, *Bacillus lautus*, *Bacillus megaterium*, *Bacillus thuringiensis*, or *Streptomyces lividans* or *Streptomyces murinus*, or gramnegative bacteria such as *E.coli*. The transformation of the bacteria may, for instance, be effected by protoplast transformation or by using competent cells in a manner known *per se*.

15           The yeast organism may favourably be selected from a species of *Saccharomyces* or *Schizosaccharomyces*, e.g. *Saccharomyces cerevisiae*. The filamentous fungus may advantageously belong to a species of *Aspergillus*, e.g. *Aspergillus oryzae* or *Aspergillus niger*. Fungal cells may be transformed by a process involving protoplast formation and transformation of the protoplasts followed by regeneration of the cell wall in a manner known *per se*. A suitable procedure for transformation of *Aspergillus* host cells is described in EP 238 023.

20           In a yet further aspect, the present invention relates to a method of producing an  $\alpha$ -amylase variant of the invention, which method comprises cultivating a host cell as described above under conditions conducive to the production of the variant and recovering the variant from the cells and/or culture medium.

25           The medium used to cultivate the cells may be any conventional medium suitable for growing the host cell in question and obtaining expression of the  $\alpha$ -amylase variant of the invention. Suitable media are available from commercial suppliers or may be prepared according to published recipes (e.g. as described in catalogues of the American Type Culture Collection).

30           The  $\alpha$ -amylase variant secreted from the host cells may conveniently be recovered from the culture medium by well-known procedures, including separating the cells

from the medium by centrifugation or filtration, and precipitating proteinaceous components of the medium by means of a salt such as ammonium sulphate, followed by the use of chromatographic procedures such as ion exchange chromatography, affinity chromatography, or the like.

5

### Industrial Applications

The  $\alpha$ -amylase variants of this invention possesses valuable properties allowing for various industrial applications. In particular the enzyme variants finds potential applications as a component in washing, dishwashing and hard surface cleaning detergent compositions, but it may also be useful in the production of sweeteners and ethanol from starch and for textile desizing. Conditions for conventional starch converting processes and liquefaction and/or saccharification processes are described in for instance US Patent No. 3,912,590 and EP patent publications Nos. 252,730 and 63,909.

10

Production of sweeteners from starch: A "traditional" process for conversion of starch to fructose syrups normally consists of three consecutive enzymatic processes, viz. a liquefaction process followed by a saccharification process and an isomerization process. During the liquefaction process, starch is degraded to dextrins by an  $\alpha$ -amylase (e.g. Termamyl™) at pH values between 5.5 and 6.2 and at temperatures of 95-160°C for a period of approx. 2h. In order to ensure an optimal enzyme stability under these conditions, 1mM of calcium is added (40 ppm free calcium ions).

15

20

After the liquefaction process the dextrins are converted into dextrose by addition of a glucoamylase (e.g. AMG™) and a debranching enzyme, such as an isoamylase or a pullulanase (e.g. Promozyme™). Before this step the pH is reduced to a value below 4.5, maintaining the high temperature (above 95°C), and the liquefying  $\alpha$ -amylase activity is denatured. The temperature is lowered to 60°C, and glucoamylase and debranching enzyme are added. The saccharification process proceeds for 24-72 hours.

25

After the saccharification process the pH is increased to a value in the range of 6-8, preferably pH 7.5, and the calcium is removed by ion exchange. The dextrose syrup is then converted into high fructose syrup using, e.g., an immobilized glucoseisomerase (such as Sweetzyme™).

30

At least 3 enzymatic improvements of this process could be obtained. All three improvements could be seen as individual benefits, but any combination (e.g. 1+2, 1+3, 2+3 or 1+2+3) could be employed:

5 Improvement 1. Reduction of the calcium dependency of the liquefying  $\alpha$ -amylase.

Addition of free calcium is required to ensure adequately high stability of the  $\alpha$ -amylase, but free calcium strongly inhibits the activity of the glucoseisomerase and needs to be removed, by means of an expensive unit operation, to an extent which reduces the level of free calcium to below 3-5 ppm. Cost savings could be obtained if such an operation could be avoided and the liquefaction process could be performed without addition of free calcium ions.

To achieve that, a less calcium-dependent Termamyl-like  $\alpha$ -amylase which is stable and highly active at low concentrations of free calcium ( $< 40$  ppm) is required. Such a Termamyl-like  $\alpha$ -amylase should have a pH optimum at a pH in the range of 4.5-6.5, preferably in the range of 4.5-5.5.

10 Improvement 2. Reduction of formation of unwanted Maillard products

The extent of formation of unwanted Maillard products during the liquefaction process is dependent on the pH. Low pH favours reduced formation of Maillard products. It would thus be desirable to be able to lower the process pH from around pH 6.0 to a value around pH 4.5; unfortunately, all commonly known, thermostable Termamyl-like  $\alpha$ -amylases are not very stable at low pH (i.e. pH  $< 6.0$ ) and their specific activity is generally low.

Achievement of the above-mentioned goal requires a Termamyl-like  $\alpha$ -amylase which is stable at low pH in the range of 4.5-5.5 and at free calcium concentrations in the range of 0-40 ppm, and which maintains a high specific activity.

25 Improvement 3.

It has been reported previously (US patent 5,234,823) that when saccharifying with *A. niger* glucoamylase and *B. acidopullulyticus* pullulanase, the presence of residual  $\alpha$ -amylase activity from the liquefaction process can lead to lower yields of dextrose if the  $\alpha$ -amylase is not inactivated before the saccharification stage. This inactivation can typically be

carried out by adjusting the pH to below 4.3 at 95°C, before lowering the temperature to 60°C for saccharification.

The reason for this negative effect on dextrose yield is not fully understood, but it is assumed that the liquefying  $\alpha$ -amylase (for example Termamyl™ 120 L from *B. licheniformis*) generates "limit dextrins" (which are poor substrates for *B. acidopullulyticus* pullulanase) by hydrolysing 1,4- $\alpha$ -glucosidic linkages close to and on both sides of the branching points in amylopectin. Hydrolysis of these limit dextrins by glucoamylase leads to a build-up of the trisaccharide panose, which is only slowly hydrolysed by glucoamylase.

The development of a thermostable  $\alpha$ -amylase which does not suffer from this disadvantage would be a significant process improvement, as no separate inactivation step would be required.

If a Termamyl-like, low-pH-stable  $\alpha$ -amylase is developed, an alteration of the specificity could be an advantage needed in combination with increased stability at low pH.

The methodology and principles of the present invention make it possible to design and produce variants according to the invention having required properties as outlined above. In this connection, mutations in a Termamyl-like  $\alpha$ -amylase [for example Termamyl itself (*B. licheniformis*  $\alpha$ -amylase; SEQ ID NO: 2); or a Termamyl-like  $\alpha$ -amylase having an N-terminal amino acid sequence (i.e. the partial sequence up to the amino acid position corresponding to position 35 in Termamyl) which is identical to that in *B. amyloliquefaciens*  $\alpha$ -amylase (SEQ ID NO: 4), i.e. a Termamyl-like  $\alpha$ -amylase having the following N-terminal sequence relative to amino acid sequence of Termamyl:

A1\*+N2\*+L3V+M15T+R23K+S29A+A30E+Y31H+A33S+E34D+H35I, where an asterisk (\*) indicates deletion of the amino acid residue in question] at positions corresponding to any of the following positions in Termamyl are particularly interesting:

H133

H156

A181

A209

G310

H450

V128

N104

V54  
S187  
H293  
A294

5 (where each of the latter amino acid residues may be replaced by any other amino acid residue, i.e. any other residue chosen among A, R, N, D, C, Q, E, G, H, I, L, K, M, F, P, S, T, W, Y and V), as well as the following triple deletions:

K370\*+G371\*+D372\*

D372\*+S373\*+Q374\*

10 Particularly preferred substitutions at the above-indicated positions are the following:

H133I

H156Y

A181T

15 A209V

G310D

H450Y

V128E

N104D

20 V54W,Y,F,I,L

S187D

H293Y

A294V.

25 Any combination of one or more (i.e. one, two, three, four..... etc.) of the above indicated mutations may appropriately be effected in a Termamyl-like  $\alpha$ -amylase in the context in question, and particularly interesting variants of the invention in the context of achieving one or more of the above-mentioned improvements in relation to the starch liquefaction behaviour of  $\alpha$ -amylases include variants comprising combinations of multiple mutations corresponding to the following combinations of mutations in Termamyl itself:

30 H133I+H156Y+A181T+A209V+G310D+H450Y+V128E+N104D+V54W+S187D+H293Y+A294V+K370\*+G371\*+D372\*;

H133I+H156Y+A181T+A209V+G310D+H450Y+V128E+N104D+V54W+S187D+H293Y+A294V+D372\*+S373\*+Q374\*;

H133I+H156Y+A181T+A209V+G310D+H450Y+V128E+N104D+V54Y+S187D+H293Y+A294V+K370\*+G371\*+D372\*;

5 H133I+H156Y+A181T+A209V+G310D+H450Y+V128E+N104D+V54Y+S187D+H293Y+A294V+D372\*+S373\*+Q374\*;

H133I+H156Y+A181T+A209V+G310D+H450Y+V128E+N104D+V54F+S187D+H293Y+A294V+K370\*+G371\*+D372\*;

10 H133I+H156Y+A181T+A209V+G310D+H450Y+V128E+N104D+V54F+S187D+H293Y+A294V+D372\*+S373\*+Q374\*;

H133I+H156Y+A181T+A209V+G310D+H450Y+V128E+N104D+V54I+S187D+H293Y+A294V+K370\*+G371\*+D372\*;

H133I+H156Y+A181T+A209V+G310D+H450Y+V128E+N104D+V54I+S187D+H293Y+A294V+D372\*+S373\*+Q374\*;

15 H133I+H156Y+A181T+A209V+G310D+H450Y+V128E+N104D+V54L+S187D+H293Y+A294V+K370\*+G371\*+D372\*;

H133I+H156Y+A181T+A209V+G310D+H450Y+V128E+N104D+V54L+S187D+H293Y+A294V+D372\*+S373\*+Q374\*;

#### 20 Detergent Compositions

According to the invention, the  $\alpha$ -amylase may typically be a component of a detergent composition. As such, it may be included in the detergent composition in the form of a non-dusting granulate, a stabilized liquid, or a protected enzyme. Non-dusting granulates may be produced, e.g. as disclosed in US 4,106,991 and 4,661,452 (both to Novo Industri A/S) and may optionally be coated by methods known in the art. Examples of waxy coating materials are poly(ethylene oxide) products (polyethyleneglycol, PEG) with mean molar weights of 1000 to 20000, ethoxylated nonylphenols having from 16 to 50 ethylene oxide units; ethoxylated fatty alcohols in which the alcohol contains from 12 to 20 carbon atoms and in which there are 15 to 80 ethylene oxide units; fatty alcohols; fatty acids; and mono- and di- and triglycerides of fatty acids. Examples of film-forming coating materials suitable for application by fluid bed techniques are given in patent GB 1483591. Liquid enzyme preparations may, for instance, be stabilized by adding a polyol such as propylene glycol, a

sugar or sugar alcohol, lactic acid or boric acid according to established methods. Other enzyme stabilizers are well known in the art. Protected enzymes may be prepared according to the method disclosed in EP 238,216.

The detergent composition of the invention may be in any convenient form, e.g. as powder, granules, paste or liquid. A liquid detergent may be aqueous, typically containing up to 70% of water and 0-30% of organic solvent, or nonaqueous.

The detergent composition comprises one or more surfactants, each of which may be anionic, nonionic, cationic, or zwitterionic. The detergent will usually contain 0-50% of anionic surfactant such as linear alkylbenzenesulfonate (LAS), alpha-olefinsulfonate (AOS), alkyl sulfate (fatty alcohol sulfate) (AS), alcohol ethoxysulfate (AEOS or AES), secondary alkanesulfonates (SAS), alpha-sulfo fatty acid methyl esters, alkyl- or alkenylsuccinic acid or soap. It may also contain 0-40% of nonionic surfactant such as alcohol ethoxylate (AEO or AE), carboxylated alcohol ethoxylates, nonylphenol ethoxylate, alkylpolyglycoside, alkyldimethylamineoxide, ethoxylated fatty acid monoethanolamide, fatty acid monoethanolamide, or polyhydroxy alkyl fatty acid amide (e.g. as described in WO 92/06154).

The detergent composition may additionally comprise one or more other enzymes, such as lipase, cutinase, protease, cellulase, peroxidase, e.g., laccase.

The detergent may contain 1-65% of a detergent builder or complexing agent such as zeolite, diphosphate, triphosphate, phosphonate, citrate, nitrilotriacetic acid (NTA), ethylenediaminetetraacetic acid (EDTA), diethylenetriaminepentaacetic acid (DTMPA), alkyl- or alkenylsuccinic acid, soluble silicates or layered silicates (e.g. SKS-6 from Hoechst). The detergent may also be unbuilt, i.e. essentially free of detergent builder.

The detergent may comprise one or more polymers. Examples are carboxymethylcellulose (CMC), poly(vinylpyrrolidone) (PVP), polyethyleneglycol (PEG), poly(vinyl alcohol) (PVA), polycarboxylates such as polyacrylates, maleic/acrylic acid copolymers and lauryl methacrylate/acrylic acid copolymers.

The detergent may contain a bleaching system which may comprise a  $H_2O_2$  source such as perborate or percarbonate which may be combined with a peracid-forming bleach activator such as tetraacetylenediamine (TAED) or nonanoyloxybenzenesulfonate (NOBS). Alternatively, the bleaching system may comprise peroxy acids of e.g. the amide, imide, or sulfone type.



The enzymes of the detergent composition of the invention may be stabilized using conventional stabilizing agents, e.g. a polyol such as propylene glycol or glycerol, a sugar or sugar alcohol, lactic acid, boric acid, or a boric acid derivative as e.g. an aromatic borate ester, and the composition may be formulated as described in e.g. WO 92/19709 and WO 92/19708.

The detergent may also contain other conventional detergent ingredients such as e.g. fabric conditioners including clays, foam boosters, suds suppressors, anti-corrosion agents, soil-suspending agents, anti-soil redeposition agents, dyes, bactericides, optical brighteners, or perfume.

The pH (measured in aqueous solution at use concentration) will usually be neutral or alkaline, e.g. 7-11.

Particular forms of detergent compositions within the scope of the invention include:

1) A detergent composition formulated as a granulate having a bulk density of at least 600 g/l comprising

Linear alkylbenzenesulfonate (calculated as acid)	7 - 12%
Alcohol ethoxysulfate (e.g., C <sub>12-18</sub> alcohol, 1-2 EO) or alkyl sulfate (e.g., C <sub>16-18</sub> )	1 - 4%
Alcohol ethoxylate (e.g., C <sub>14-15</sub> alcohol, 7 EO)	5 - 9%
Sodium carbonate (as Na <sub>2</sub> CO <sub>3</sub> )	14 - 20%
Soluble silicate (as Na <sub>2</sub> O, 2SiO <sub>2</sub> )	2 - 6%
Zeolite (as NaAlSiO <sub>4</sub> )	15 - 22%
Sodium sulfate (as Na <sub>2</sub> SO <sub>4</sub> )	0 - 6%
Sodium citrate/citric acid (as C <sub>6</sub> H <sub>5</sub> Na <sub>3</sub> O <sub>7</sub> /C <sub>6</sub> H <sub>8</sub> O <sub>7</sub> )	0 - 15%
Sodium perborate (as NaBO <sub>3</sub> ·H <sub>2</sub> O)	11 - 18%
TAED	2 - 6%
Carboxymethylcellulose	0 - 2%
Polymers (e.g., maleic/acrylic acid copolymer, PVP, PEG)	0 - 3%
Enzymes (calculated as pure enzyme protein)	0.0001 - 0.1%
Minor ingredients (e.g., suds suppressors, perfume, optical brightener, photobleach)	0 - 5%

2) A detergent composition formulated as a granulate having a bulk density of at least 600 g/l comprising

Linear alkylbenzenesulfonate (calculated as acid)	6 - 11%
Alcohol ethoxysulfate ( <i>e.g.</i> , C <sub>12-18</sub> alcohol, 1-2 EO or alkyl sulfate ( <i>e.g.</i> , C <sub>16-18</sub> ))	1 - 3%
Alcohol ethoxylate ( <i>e.g.</i> , C <sub>14-15</sub> alcohol, 7 EO)	5 - 9%
Sodium carbonate (as Na <sub>2</sub> CO <sub>3</sub> )	15 - 21%
Soluble silicate (as Na <sub>2</sub> O, 2SiO <sub>2</sub> )	1 - 4%
Zeolite (as NaAlSiO <sub>4</sub> )	24 - 34%
Sodium sulfate (as Na <sub>2</sub> SO <sub>4</sub> )	4 - 10%
Sodium citrate/citric acid (as C <sub>6</sub> H <sub>5</sub> Na <sub>3</sub> O <sub>7</sub> /C <sub>6</sub> H <sub>8</sub> O <sub>7</sub> )	0 - 15%
Carboxymethylcellulose	0 - 2%
Polymers ( <i>e.g.</i> , maleic/acrylic acid copolymer, PVP, PEG)	1 - 6%
Enzymes (calculated as pure enzyme protein)	0.0001 - 0.1%
Minor ingredients ( <i>e.g.</i> , suds suppressors, perfume)	0 - 5%

3) A detergent composition formulated as a granulate having a bulk density of at least 600 g/l comprising

Linear alkylbenzenesulfonate (calculated as acid)	5 - 9%
Alcohol ethoxylate ( <i>e.g.</i> , C <sub>12-15</sub> alcohol, 7 EO)	7 - 14%
Soap as fatty acid ( <i>e.g.</i> , C <sub>16-22</sub> fatty acid)	1 - 3%
Sodium carbonate (as Na <sub>2</sub> CO <sub>3</sub> )	10 - 17%
Soluble silicate (as Na <sub>2</sub> O, 2SiO <sub>2</sub> )	3 - 9%
Zeolite (as NaAlSiO <sub>4</sub> )	23 - 33%
Sodium sulfate (as Na <sub>2</sub> SO <sub>4</sub> )	0 - 4%
Sodium perborate (as NaBO <sub>3</sub> ·H <sub>2</sub> O)	8 - 16%
TAED	2 - 8%
Phosphonate ( <i>e.g.</i> , EDTMPA)	0 - 1%
Carboxymethylcellulose	0 - 2%
Polymers ( <i>e.g.</i> , maleic/acrylic acid copolymer, PVP, PEG)	0 - 3%
Enzymes (calculated as pure enzyme protein)	0.0001 - 0.1%
Minor ingredients ( <i>e.g.</i> , suds suppressors, perfume, optical brightener)	0 - 5%

4) A detergent composition formulated as a granulate having a bulk density of at least 600 g/l comprising

Linear alkylbenzenesulfonate (calculated as acid)	8 - 12%
Alcohol ethoxylate ( <i>e.g.</i> , C <sub>12-15</sub> alcohol, 7 EO)	10 - 25%
Sodium carbonate (as Na <sub>2</sub> CO <sub>3</sub> )	14 - 22%
Soluble silicate (as Na <sub>2</sub> O, 2SiO <sub>2</sub> )	1 - 5%
Zeolite (as NaAlSiO <sub>4</sub> )	25 - 35%
Sodium sulfate (as Na <sub>2</sub> SO <sub>4</sub> )	0 - 10%
Carboxymethylcellulose	0 - 2%
Polymers ( <i>e.g.</i> , maleic/acrylic acid copolymer, PVP, PEG)	1 - 3%
Enzymes (calculated as pure enzyme protein)	0.0001 - 0.1%
Minor ingredients ( <i>e.g.</i> , suds suppressors, perfume)	0 - 5%

5) An aqueous liquid detergent composition comprising

Linear alkylbenzenesulfonate (calculated as acid)	15 - 21%
Alcohol ethoxylate ( <i>e.g.</i> , C <sub>12-15</sub> alcohol, 7 EO or C <sub>12-15</sub> alcohol, 5 EO)	12 - 18%
Soap as fatty acid ( <i>e.g.</i> , oleic acid)	3 - 13%
Alkenylsuccinic acid (C <sub>12-14</sub> )	0 - 13%
Aminoethanol	8 - 18%
Citric acid	2 - 8%
Phosphonate	0 - 3%
Polymers ( <i>e.g.</i> , PVP, PEG)	0 - 3%
Borate (as B <sub>4</sub> O <sub>7</sub> )	0 - 2%
Ethanol	0 - 3%
Propylene glycol	8 - 14%
Enzymes (calculated as pure enzyme protein)	0.0001 - 0.1%
Minor ingredients ( <i>e.g.</i> , dispersants, suds suppressors, perfume, optical brightener)	0 - 5%

6) An aqueous structured liquid detergent composition comprising

Linear alkylbenzenesulfonate (calculated as acid)	15 - 21%
Alcohol ethoxylate ( <i>e.g.</i> , C <sub>12-15</sub> alcohol, 7 EO, or C <sub>12-15</sub> alcohol, 5 EO)	3 - 9%
Soap as fatty acid ( <i>e.g.</i> , oleic acid)	3 - 10%
Zeolite (as NaAlSiO <sub>4</sub> )	14 - 22%
Potassium citrate	9 - 18%
Borate (as B <sub>4</sub> O <sub>7</sub> )	0 - 2%
Carboxymethylcellulose	0 - 2%
Polymers ( <i>e.g.</i> , PEG, PVP)	0 - 3%
Anchoring polymers such as, <i>e.g.</i> , lauryl methacrylate/acrylic acid copolymer; molar ratio 25:1; MW 3800	0 - 3%
Glycerol	0 - 5%
Enzymes (calculated as pure enzyme protein)	0.0001 - 0.1%
Minor ingredients ( <i>e.g.</i> , dispersants, suds suppressors, perfume, optical brighteners)	0 - 5%

7) A detergent composition formulated as a granulate having a bulk density of at least 600 g/l comprising

Fatty alcohol sulfate	5 - 10%
Ethoxylated fatty acid monoethanolamide.	3 - 9%
Soap as fatty acid	0 - 3%
Sodium carbonate (as $\text{Na}_2\text{CO}_3$ )	5 - 10%
Soluble silicate (as $\text{Na}_2\text{O}$ , $2\text{SiO}_2$ )	1 - 4%
Zeolite (as $\text{NaAlSiO}_4$ )	20 - 40%
Sodium sulfate (as $\text{Na}_2\text{SO}_4$ )	2 - 8%
Sodium perborate (as $\text{NaBO}_3 \cdot \text{H}_2\text{O}$ )	12 - 18%
TAED	2 - 7%
Polymers (e.g., maleic/acrylic acid copolymer, PEG)	1 - 5%
Enzymes (calculated as pure enzyme protein)	0.0001 - 0.1%
Minor ingredients (e.g., optical brightener, suds suppressors, perfume)	0 - 5%

8) A detergent composition formulated as a granulate comprising

Linear alkylbenzenesulfonate (calculated as acid)	8 - 14%
Ethoxylated fatty acid monoethanolamide	5 - 11%
Soap as fatty acid	0 - 3%
Sodium carbonate (as $\text{Na}_2\text{CO}_3$ )	4 - 10%
Soluble silicate (as $\text{Na}_2\text{O}$ , $2\text{SiO}_2$ )	1 - 4%
Zeolite (as $\text{NaAlSiO}_4$ )	30 - 50%
Sodium sulfate (as $\text{Na}_2\text{SO}_4$ )	3 - 11%
Sodium citrate (as $\text{C}_6\text{H}_5\text{Na}_3\text{O}_7$ )	5 - 12%
Polymers (e.g., PVP, maleic/acrylic acid copolymer, PEG)	1 - 5%
Enzymes (calculated as pure enzyme protein)	0.0001 - 0.1%
Minor ingredients (e.g., suds suppressors, perfume)	0 - 5%

9) A detergent composition formulated as a granulate comprising

Linear alkylbenzenesulfonate (calculated as acid)	6 - 12%
Nonionic surfactant	1 - 4%
Soap as fatty acid	2 - 6%
Sodium carbonate (as $\text{Na}_2\text{CO}_3$ )	14 - 22%
Zeolite (as $\text{NaAlSiO}_4$ )	18 - 32%
Sodium sulfate (as $\text{Na}_2\text{SO}_4$ )	5 - 20%
Sodium citrate (as $\text{C}_6\text{H}_5\text{Na}_3\text{O}_7$ )	3 - 8%
Sodium perborate (as $\text{NaBO}_3 \cdot \text{H}_2\text{O}$ )	4 - 9%
Bleach activator ( <i>e.g.</i> , NOBS or TAED)	1 - 5%
Carboxymethylcellulose	0 - 2%
Polymers ( <i>e.g.</i> , polycarboxylate or PEG)	1 - 5%
Enzymes (calculated as pure enzyme protein)	0.0001 - 0.1%
Minor ingredients ( <i>e.g.</i> , optical brightener, perfume)	0 - 5%

10) An aqueous liquid detergent composition comprising

Linear alkylbenzenesulfonate (calculated as acid)	15 - 23%
Alcohol ethoxysulfate ( <i>e.g.</i> , $\text{C}_{12-15}$ alcohol, 2-3 EO)	8 - 15%
Alcohol ethoxylate ( <i>e.g.</i> , $\text{C}_{12-15}$ alcohol, 7 EO, or $\text{C}_{12-15}$ alcohol, 5 EO)	3 - 9%
Soap as fatty acid ( <i>e.g.</i> , lauric acid)	0 - 3%
Aminoethanol	1 - 5%
Sodium citrate	5 - 10%
Hydrotrope ( <i>e.g.</i> , sodium toluenesulfonate)	2 - 6%
Borate (as $\text{B}_4\text{O}_7$ )	0 - 2%
Carboxymethylcellulose	0 - 1%
Ethanol	1 - 3%
Propylene glycol	2 - 5%
Enzymes (calculated as pure enzyme protein)	0.0001 - 0.1%
Minor ingredients ( <i>e.g.</i> , polymers, dispersants, perfume, optical brighteners)	0 - 5%

11) An aqueous liquid detergent composition comprising

Linear alkylbenzenesulfonate (calculated as acid)	20 - 32%
Alcohol ethoxylate ( <i>e.g.</i> , C <sub>12-15</sub> alcohol, 7 EO, or C <sub>12-15</sub> alcohol, 5 EO)	6 - 12%
Aminoethanol	2 - 6%
Citric acid	8 - 14%
Borate (as B <sub>4</sub> O <sub>7</sub> )	1 - 3%
Polymer ( <i>e.g.</i> , maleic/acrylic acid copolymer, anchoring polymer such as, <i>e.g.</i> , lauryl methacrylate/acrylic acid copolymer)	0 - 3%
Glycerol	3 - 8%
Enzymes (calculated as pure enzyme protein)	0.0001 - 0.1%
Minor ingredients ( <i>e.g.</i> , hydrotropes, dispersants, perfume, optical brighteners)	0 - 5%

12) A detergent composition formulated as a granulate having a bulk density of at least 600 g/l comprising

Anionic surfactant (linear alkylbenzenesulfonate, alkyl sulfate, alpha-olefinsulfonate, alpha-sulfo fatty acid methyl esters, alkanesulfonates, soap)	25 - 40%
Nonionic surfactant ( <i>e.g.</i> , alcohol ethoxylate)	1 - 10%
Sodium carbonate (as Na <sub>2</sub> CO <sub>3</sub> )	8 - 25%
Soluble silicates (as Na <sub>2</sub> O, 2SiO <sub>2</sub> )	5 - 15%
Sodium sulfate (as Na <sub>2</sub> SO <sub>4</sub> )	0 - 5%
Zeolite (as NaAlSiO <sub>4</sub> )	15 - 28%
Sodium perborate (as NaBO <sub>3</sub> ·4H <sub>2</sub> O)	0 - 20%
Bleach activator (TAED or NOBS)	0 - 5%
Enzymes (calculated as pure enzyme protein)	0.0001 - 0.1%
Minor ingredients ( <i>e.g.</i> , perfume, optical brighteners)	0 - 3%

13) Detergent compositions as described in compositions 1-12 wherein all or part of the linear alkylbenzenesulfonate is replaced by (C<sub>12</sub>-C<sub>18</sub>) alkyl sulfate.

14) A detergent composition formulated as a granulate having a bulk density of at least 600 g/l comprising

(C <sub>12</sub> -C <sub>18</sub> ) alkyl sulfate	9 - 15%
Alcohol ethoxylate	3 - 6%
Polyhydroxy alkyl fatty acid amide	1 - 5%
Zeolite (as NaAlSiO <sub>4</sub> )	10 - 20%
Layered disilicate (e.g., SK56 from Hoechst)	10 - 20%
Sodium carbonate (as Na <sub>2</sub> CO <sub>3</sub> )	3 - 12%
Soluble silicate (as Na <sub>2</sub> O, 2SiO <sub>2</sub> )	0 - 6%
Sodium citrate	4 - 8%
Sodium percarbonate	13 - 22%
TAED	3 - 8%
Polymers (e.g., polycarboxylates and PVP)	0 - 5%
Enzymes (calculated as pure enzyme protein)	0.0001 - 0.1%
Minor ingredients (e.g., optical brightener, photo bleach, perfume, suds suppressors)	0 - 5%

15) A detergent composition formulated as a granulate having a bulk density of at least 600 g/l comprising

(C <sub>12</sub> -C <sub>18</sub> ) alkyl sulfate	4 - 8%
Alcohol ethoxylate	11 - 15%
Soap	1 - 4%
Zeolite MAP or zeolite A	35 - 45%
Sodium carbonate (as Na <sub>2</sub> CO <sub>3</sub> )	2 - 8%
Soluble silicate (as Na <sub>2</sub> O, 2SiO <sub>2</sub> )	0 - 4%
Sodium percarbonate	13 - 22%
TAED	1 - 8%
Carboxymethyl cellulose	0 - 3%
Polymers (e.g., polycarboxylates and PVP)	0 - 3%
Enzymes (calculated as pure enzyme protein)	0.0001 - 0.1%
Minor ingredients (e.g., optical brightener, phosphonate, perfume)	0 - 3%



16) Detergent formulations as described in 1) - 15) which contain a stabilized or encapsulated peracid, either as an additional component or as a substitute for already specified bleach systems.

17) Detergent compositions as described in 1), 3), 7), 9) and 12) wherein perborate is replaced by percarbonate.

18) Detergent compositions as described in 1), 3), 7), 9), 12), 14) and 15) which additionally contain a manganese catalyst. The manganese catalyst may, e.g., be one of the compounds described in "Efficient manganese catalysts for low-temperature bleaching", Nature 369, 1994, pp. 637-639.

19) Detergent composition formulated as a nonaqueous detergent liquid comprising a liquid nonionic surfactant such as, e.g., linear alkoxyated primary alcohol, a builder system (e.g. phosphate), enzyme and alkali. The detergent may also comprise anionic surfactant and/or a bleach system.

The  $\alpha$ -amylase variant of the invention may be incorporated in concentrations conventionally employed in detergents. It is at present contemplated that, in the detergent composition of the invention, the  $\alpha$ -amylase may be added in an amount corresponding to 0.00001-1 mg (calculated as pure enzyme protein) of  $\alpha$ -amylase per liter of wash liquor.

#### Dishwashing Composition

The dishwashing detergent composition comprises a surfactant which may be anionic, non-ionic, cationic, amphoteric or a mixture of these types. The detergent will contain 0-90% of non-ionic surfactant such as low- to non-foaming ethoxylated propoxylated straight-chain alcohols.

The detergent composition may contain detergent builder salts of inorganic and/or organic types. The detergent builders may be subdivided into phosphorus-containing and non-phosphorus-containing types. The detergent composition usually contains 1-90% of detergent builders.

Examples of phosphorus-containing inorganic alkaline detergent builders, when present, include the water-soluble salts especially alkali metal pyrophosphates,

orthophosphates, and polyphosphates. An example of phosphorus-containing organic alkaline detergent builder, when present, includes the water-soluble salts of phosphonates. Examples of non-phosphorus-containing inorganic builders, when present, include water-soluble alkali metal carbonates, borates and silicates as well as the various types of water-insoluble crystalline or amorphous aluminosilicates of which zeolites are the best-known representatives.

Examples of suitable organic builders include the alkali metal, ammonium and substituted ammonium, citrates, succinates, malonates, fatty acid sulphonates, carboxymethoxy succinates, ammonium polyacetates, carboxylates, polycarboxylates, aminopolycarboxylates, polyacetyl carboxylates and polyhydroxysulphonates.

Other suitable organic builders include the higher molecular weight polymers and co-polymers known to have builder properties, for example appropriate polyacrylic acid, polymaleic and polyacrylic/polymaleic acid copolymers and their salts.

The dishwashing detergent composition may contain bleaching agents of the chlorine/bromine-type or the oxygen-type. Examples of inorganic chlorine/bromine-type bleaches are lithium, sodium or calcium hypochlorite and hypobromite as well as chlorinated trisodium phosphate. Examples of organic chlorine/bromine-type bleaches are heterocyclic N-bromo and N-chloro imides such as trichloroisocyanuric, tribromoisocyanuric, dibromoisocyanuric and dichloroisocyanuric acids, and salts thereof with water-solubilizing cations such as potassium and sodium. Hydantoin compounds are also suitable.

The oxygen bleaches are preferred, for example in the form of an inorganic persalt, preferably with a bleach precursor or as a peroxy acid compound. Typical examples of suitable peroxy bleach compounds are alkali metal perborates, both tetrahydrates and monohydrates, alkali metal percarbonates, persilicates and perphosphates. Preferred activator materials are TAED and glycerol triacetate.

The dishwashing detergent composition of the invention may be stabilized using conventional stabilizing agents for the enzyme(s), e.g. a polyol such as e.g. propylene glycol, a sugar or a sugar alcohol, lactic acid, boric acid, or a boric acid derivative, e.g. an aromatic borate ester.

The dishwashing detergent composition of the invention may also contain other conventional detergent ingredients, e.g. deflocculant material, filler material, foam depressors,

anti-corrosion agents, soil-suspending agents, sequestering agents, anti-soil redeposition agents, dehydrating agents, dyes, bactericides, fluorescers, thickeners and perfumes.

Finally, the  $\alpha$ -amylase variant of the invention may be used in conventional dishwashing detergents, e.g. in any of the detergents described in any of the following patent publications:

EP 518719, EP 518720, EP 518721, EP 516553, EP 516554, EP 516555, GB 2200132, DE 3741617, DE 3727911, DE 4212166, DE 4137470, DE 3833047, WO 93/17089, DE 4205071, WO 52/09680, WO 93/18129, WO 93/04153, WO 92/06157, WO 92/08777, EP 429124, WO 93/21299, US 5141664, EP 561452, EP 561446, GB 2234980, WO 93/03129, EP 481547, EP 530870, EP 533239, EP 554943, EP 346137, US 5112518, EP 318204, EP 318279, EP 271155, EP 271156, EP 346136, GB 2228945, CA 2006687, WO 93/25651, EP 530635, EP 414197, US 5240632.

## EXAMPLES

### EXAMPLE 1

#### Example on Homology building of TERM

The overall homology of the *B. licheniformis*  $\alpha$ -amylase (in the following referred to as TERM) to other Termamyl-like  $\alpha$ -amylases is high and the percent similarity is extremely high. The similarity calculated for TERM to BSG (the *B. stearothermophilus*  $\alpha$ -amylase with SEQ ID NO: 6), and BAN (the *B. amyloliquefaciens*  $\alpha$ -amylase with SEQ ID NO: 4) using the University of Wisconsin Genetics Computer Group's program GCG gave 89% and 78%, respectively. TERM has a deletion of 2 residues between residue G180 and K181 compared to BAN and BSG. BSG has a deletion of 3 residues between G371 and I372 in comparison with BAN and TERM. Further BSG has a C-terminal extension of more than 20 residues compared to BAN and TERM. BAN has 2 residues less and TERM has one residue less in the N-terminal compared to BSG.

The structure of the *B. licheniformis* (TERM) and of the *B. amyloliquefaciens*  $\alpha$ -amylase (BAN), respectively, was model built on the structure disclosed in Appendix 1 herein. The structure of other Termamyl-like  $\alpha$ -amylases (e.g. those disclosed herein) may be built analogously.

In comparison with the  $\alpha$ -amylase used for elucidating the present structure, TERM differs in that it lacks two residues around 178-182. In order to compensate for this

in the model structure, the HOMOLOGY program from BIOSYM was used to substitute the residues in equivalent positions in the structure (not only structurally conserved regions) except for the deletion point. A peptide bond was established between G179(G177) and K180(K180) in TERM(BAN). The close structural relationship between the solved structure and the model structure (and thus the validity of the latter) is indicated by the presence of only very few atoms found to be too close together in the model.

To this very rough structure of TERM was then added all waters (605) and ions (4 Calcium and 1 Sodium) from the solved structure (Appendix 1) at the same coordinates as for said solved structure using the INSIGHT program. This could be done with only few overlaps - in other words with a very nice fit. This model structure were then minimized using 200 steps of Steepest descent and 600 steps of Conjugated gradient (see Brooks et al 1983, J. Computational Chemistry 4, p.187-217). The minimized structure was then subjected to molecular dynamics, 5ps heating followed by up to 200ps equilibration but more than 35ps. The dynamics as run with the Verlet algorithm and the equilibration temperature 300K were kept using the Berendsen coupling to a waterbath (Berendsen et. al., 1984, J. Chemical Physics 81, p. 3684-3690). Rotations and translations were removed every picosecond. The potential energy became stable after appr. 35ps equilibration. A mean dynamics structure was extracted and can be used for further analysis.

## EXAMPLE 2

### Determination of residues within 10Å from the ions present in the solved structure

The coordinates of Appendix 1 are read into the INSIGHT program provided by BIOSYM technologies. The spatial coordinates are presented showing the bonds between the atoms. The ions are presented as well as the water atoms. The program package part of creating subset are used to create a 10Å subset around the Calcium and the Sodium ions in the structure using the command ZONE. All residues having an atom within the 10Å are compiled and written out by the LIST MOLECULE command. By giving the ions the name ium in the coordinate file a 10Å sphere around all atoms called ium is compiled. The specific residues identified in this manner are given further above in the section entitled "Ca<sup>2+</sup> dependency".

### EXAMPLE 3

#### Determination of cavities in the solved structure (Appendix 1)

The solved structure exhibits many internal holes and cavities. When analysing for such cavities the Connolly program is normally used (Lee, B. and Richards, F.M. (1971) J. Mol. Biol. 55,p. 379-400). The program uses a probe with radius to search the external and internal surface of the protein. The smallest hole observable in this way has the probe radius.

To analyse the solved structure a modified version of the Connolly program included in the program of INSIGHT were used. First the water molecules and the ions were removed by unmerging these atoms from the solved structure. By using the command MOLECULE SURFACE SOLVENT the solvent accessible surface area were calculated for all atoms and residues using a probe radius of 1.4Å, and displayed on the graphics screen together with the model of the solved structure. The internal cavities where then seen as dot surfaces with no connections to external surface.

Mutant suggestions for filling out the holes are given in the specification (in the section entitled "Variants with increased thermostability and/or altered temperature optimum"). By using the homology build structures or/and the sequence alignment mutations for the homologous structures of TERM and BSG and BAN can be made.

### EXAMPLE 4

#### Construction of Termamyl™ variants in accordance with the invention

Termamyl (SEQ ID NO: 2) is expressed in *B. subtilis* from a plasmid denoted pDN1528. This plasmid contains the complete gene encoding Termamyl, *amyL*, the expression of which is directed by its own promoter. Further, the plasmid contains the origin of replication, *ori*, from plasmid pUB110 and the *cat* gene from plasmid pC194 conferring resistance towards chloramphenicol. pDN1528 is shown in Fig. 9.

A specific mutagenesis vector containing a major part of the coding region of SEQ ID NO: 1 was prepared. The important features of this vector, denoted pJeEN1, include an origin of replication derived from the pUC plasmids, the *cat* gene conferring resistance towards chloramphenicol, and a frameshift-containing version of the *bla* gene, the wild type of which normally confers resistance towards ampicillin (amp<sup>R</sup> phenotype). This mutated version results in an amp<sup>S</sup> phenotype. The plasmid pJeEN1 is shown in Fig. 10, and the *E.*

*coli* origin of replication, *ori*, *bla*, *cat*, the 5'-truncated version of the Termamyl amylase gene, and selected restriction sites are indicated on the plasmid.

Mutations are introduced in *amyL* by the method described by Deng and Nickoloff (1992, *Anal. Biochem.* 200, pp. 81-88) except that plasmids with the "selection primer" (primer #6616; see below) incorporated are selected based on the *amp*<sup>R</sup> phenotype of transformed *E. coli* cells harboring a plasmid with a repaired *bla* gene, instead of employing the selection by restriction enzyme digestion outlined by Deng and Nickoloff. Chemicals and enzymes used for the mutagenesis were obtained from the Chameleon™ mutagenesis kit from Stratagene (catalogue number 200509).

After verification of the DNA sequence in variant plasmids, the truncated gene, containing the desired alteration, is subcloned into pDN1528 as a *Pst*I-*Eco*RI fragment and transformed into a protease- and amylase-depleted *Bacillus subtilis* strain in order to express the variant enzyme.

The Termamyl variant V54W was constructed by the use of the following mutagenesis primer (written 5' to 3', left to right):

PG GTC GTA GGC ACC GTA GCC CCA ATC CGC TTG

The Termamyl variant A52W + V54W was constructed by the use of the following mutagenesis primer (written 5' to 3', left to right):

PG GTC GTA GGC ACC GTA GCC CCA ATC CCA TTG GCT CG

Primer #6616 (written 5' to 3', left to right; P denotes a 5' phosphate):

P CTG TGA CTG GTG AGT ACT CAA CCA AGT C

## EXAMPLE 5

### Saccharification in the presence of "residual" $\alpha$ -amylase activity

Two appropriate Termamyl variants with altered specificity were evaluated by saccharifying a DE 10 (DE = dextrose equivalent) maltodextrin substrate with *A. niger* glucoamylase and *B. acidopullulyticus* pullulanase under conditions where the variant amylase was active.

Saccharification: Substrates for saccharification were prepared by dissolving 230 g DE 10 spray-dried maltodextrin, prepared from common corn starch, in 460 ml boiling deionized water and adjusting the dry substance (DS) content to approximately 30% w/w. The pH was

adjusted to 4.7 (measured at 60°C) and aliquots of substrate corresponding to 15 g dry weight were transferred to 50 ml blue cap glass flasks.

The flasks were then placed in a shaking water bath equilibrated at 60°C, and the enzymes added. The pH was readjusted to 4.7 where necessary.

The following enzymes were used:

Glucoamylase: AMG™ (Novo Nordisk A/S); dosage 0.18 AG/g DS

Pullulanase: Promozyme™ (Novo Nordisk A/S);  
dosage 0.06 PUN/g DS

α-Amylases: Termamyl™ (Novo Nordisk A/S); dosage 60 NU/g DS

Termamyl variant V54W; dosage 60 NU/g DS

Termamyl variant V54W + A52W; dosage 60 NU/g DS

2 ml samples were taken periodically. The pH of each sample was adjusted to about 3.0, and the sample was then heated in a boiling water bath for 15 minutes to inactivate the enzymes. After cooling, the samples were treated with approximately 0.1 g mixed-bed ion exchange resin (BIO-Rad 501-X (D)) for 30 minutes on a rotary mixer and then filtered. The carbohydrate composition of each sample was determined by HPLC. The following results were obtained after 72 hours [DP<sub>n</sub> denotes a dextrose (D-glucose) oligomer with n glucose units]:

α-amylase	%DP <sub>1</sub>	%DP <sub>2</sub>	%DP <sub>3</sub>	%DP <sub>4</sub>
None (control)	95.9	2.8	0.4	1.0
V54W	96.0	2.9	0.4	0.8
V54W + A52W	95.9	2.8	0.4	0.8
Termamyl™	95.6	2.8	0.8	0.8

It can be seen from the above results that compared with the control (no α-amylase activity present during liquefaction), the presence of α-amylase activity from variants V54W and V54W + A52W did not lead to elevated panose (DP<sub>3</sub>) levels. In contrast, Termamyl α-amylase activity resulted in higher levels of panose and a subsequent loss of D-glucose (DP<sub>1</sub>) yield.

Thus, if  $\alpha$ -amylase variants V54W or V54W + A52W are used for starch liquefaction, it will not be necessary to inactivate the residual  $\alpha$ -amylase activity before the commencement of saccharification.

## EXAMPLE 6

### Calcium-binding affinity of $\alpha$ -amylase variants of the invention


Unfolding of amylases by exposure to heat or to denaturants such as guanidine hydrochloride is accompanied by a decrease in fluorescence. Loss of calcium ions leads to unfolding, and the affinity of  $\alpha$ -amylases for calcium can be measured by fluorescence measurements before and after incubation of each  $\alpha$ -amylase (e.g. at a concentration of 10  $\mu$ g/ml) in a buffer (e.g. 50 mM HEPES, pH 7) with different concentrations of calcium (e.g. in the range of 1  $\mu$ M-100 mM) or of EGTA (e.g. in the range of 1-1000  $\mu$ M) [EGTA = 1,2-di(2-aminoethoxy)ethane-*N,N,N',N'*-tetraacetic acid] for a sufficiently long period of time (such as 22 hours at 55°C).

The measured fluorescence  $F$  is composed of contributions from the folded and unfolded forms of the enzyme. The following equation can be derived to describe the dependence of  $F$  on calcium concentration ( $[Ca]$ ):

$$F = [Ca]/(K_{diss} + [Ca])(\alpha_N - \beta_N \log([Ca])) +$$

$$K_{diss}/(K_{diss} + [Ca])(\alpha_U - \beta_U \log([Ca]))$$

where  $\alpha_N$  is the fluorescence of the native (folded) form of the enzyme,  $\beta_N$  is the linear dependence of  $\alpha_N$  on the logarithm of the calcium concentration (as observed experimentally),  $\alpha_U$  is the fluorescence of the unfolded form and  $\beta_U$  is the linear dependence of  $\alpha_U$  on the logarithm of the calcium concentration.  $K_{diss}$  is the apparent calcium-binding constant for an equilibrium process as follows:



In fact, unfolding proceeds extremely slowly and is irreversible. The rate of unfolding is dependent on calcium concentration, and the dependency for a given  $\alpha$ -amylase provides a measure of the Ca-binding affinity of the enzyme. By defining a standard set of reaction conditions (e.g. 22 hours at 55°C), a meaningful comparison of  $K_{diss}$  for different  $\alpha$ -amylases can be made. The calcium dissociation curves for  $\alpha$ -amylases in general can be fitted to the equation above, allowing determination of the corresponding values of  $K_{diss}$ .



The following values for  $K_{\text{diss}}$  were obtained for a parent Termamyl-like  $\alpha$ -amylase having the amino acid sequence shown in SEQ ID NO: 1 of WO 95/26397 and for the indicated variant thereof according to the invention:

5	$\alpha$ -Amylase	$K_{\text{diss}}$ (mol/l)
	L351C + M430C + T183* + G184*	$1.7 (\pm 0.5) \times 10^{-3}$
	Parent	$3.5 (\pm 1.1) \times 10^{-1}$

10 It is apparent from the above that the calcium-binding affinity of the variant in question binds calcium significantly more strongly than the parent, and thereby has a correspondingly lower calcium dependency than the parent.

## REFERENCES CITED

- Klein, C., et al., *Biochemistry* 1992, **31**, 8740-8746,
- Mizuno, H., et al., *J. Mol. Biol.* (1993) **234**, 1282-1283,
- Chang, C., et al, *J. Mol. Biol.* (1993) **229**, 235-238,
- 5 Larson, S.B., *J. Mol. Biol.* (1994) **235**, 1560-1584,
- Lawson, C.L., *J. Mol. Biol.* (1994) **236**, 590-600,
- Qian, M., et al., *J. Mol. Biol.* (1993) **231**, 785-799,
- Brady, R.L., et al., *Acta Crystallogr. sect. B*, **47**, 527-535,
- Swift, H.J., et al., *Acta Crystallogr. sect. B*, **47**, 535-544
- 10 A. Kadziola, Ph.D. Thesis: "An alpha-amylase from Barley and its Complex with a Substrate Analogue Inhibitor Studied by X-ray Crystallography", Department of Chemistry University of Copenhagen 1993
- MacGregor, E.A., *Food Hydrocolloids*, 1987, Vol.1, No. 5-6, p.
- B. Diderichsen and L. Christiansen, Cloning of a maltogenic  $\alpha$ -amylase from *Bacillus*  
 5 *stearothermophilus*, FEMS Microbiol. letters: 56: pp. 53-60 (1988)
- Hudson et al., *Practical Immunology*, Third edition (1989), Blackwell Scientific Publications,
- Sambrook et al., Molecular Cloning: A Laboratory Manual, 2nd Ed., Cold Spring Harbor, 1989
- S.L. Beaucage and M.H. Caruthers, Tetrahedron Letters **22**, 1981, pp. 1859-1869
- 20 Matthes et al., The EMBO J. **3**, 1984, pp. 801-805.
- R.K. Saiki et al., Science **239**, 1988, pp. 487-491.
- Morinaga et al., (1984, *Biotechnology* 2:646-639)
- Nelson and Long, Analytical Biochemistry **180**, 1989, pp. 147-151
- Hunkapiller et al., 1984, *Nature* 310:105-111
- 25 R. Higuchi, B. Krummel, and R.K. Saiki (1988). A general method of *in vitro* preparation and specific mutagenesis of DNA fragments: study of protein and DNA interactions. *Nucl. Acids Res.* **16**:7351-7367.
- Dubnau et al., 1971, J. Mol. Biol. **56**, pp. 209-221.
- Gryczan et al., 1978, J. Bacteriol. **134**, pp. 318-329.
- 30 S.D. Erlich, 1977, Proc. Natl. Acad. Sci. **74**, pp. 1680-1682.
- Boel et al., 1990, Biochemistry **29**, pp. 6244-6249.

## SEQUENCE LISTING

In the following SEQ ID NO: 1, 3, 5 the 5', coding sequence and 3' sequence of the relevant  $\alpha$ -amylase genes are illustrated. The 5' sequence is the first separate part of the sequence written with lower case letters, the coding sequence is the intermediate part of the sequence, where the signal sequence is written with lower case letters and the sequence encoding the mature  $\alpha$ -amylase is written with upper case letters, and the 3' sequence is the third separate part of the sequence written with lower case letters.

### SEQ ID NO: 1

cggaagattggaagtacaaaaataagcaaaagattgtcaatcatgtcatgagccatgcgg-gagacggaaaaatcgtctta  
atgcacgatatttatgcaacgttcgcagatgctgctgaa-gagattattaaaaagctgaaagcaaaaggctatcaattggt  
aactgtatctcagcttgaagaagtgaagaagcagagaggctattgaataaatgagtagaagcgccatatcggcgcttttc  
tttgaagaaaaatagggaaaaatggtacttgttaaaattcggaatattatacaacatcatatgtttcacattgaaagggaggagaatc  
atgaaacaacaaaaacggcctttacgcccgattgctgacgctgttatttgcgctcatcttcttgcctgc  
ctcattctgcagcagcggcgGCAAATCTTAATGGGACGCTGATGCAGTATTTTGAATGGTAC  
ATGCCCAATGACGGCCAACATTGGAGGCGTTTGCAAACGACTCGGCATATTTGG  
CTGAACACGGTATTACTGCCGTCTGGATTCCCCCGGCATATAAGGGAACGAGCCA  
AGCGGATGTGGGCTACGGTGCTTACGACCTTTATGATTTAGGGGAGTTTCATCAAA  
AAGGGACGGTTCGGACAAAGTACGGCACAAAAGGAGAGCTGCAATCTGCGATCA  
AAAGTCTTCATTCCCGCGACATTAACGTTTACGGGGATGTGGTCATCAACCACAA  
AGGCGGCGCTGATGCGACCGAAGATGTAACCGCGGTTGAAGTCGATCCCGCTGA  
CCGCAACCG CGTAATTTCAAGGAGAACACCTAATTAAAGCCTGGACACATTTTCAT  
TTTCCGGGGCGCGGCAGCACATACAGCGATTTTA AATGGCATTGGTACCATTTTGA  
CGGAACCGATTGGGACGAGTCCCGAAAGCTGAACCGCATCTATAAGTTTCAAGG  
AAAGGCTTGGGATTGGGAAGTTTCCAATGAAAACGGCAACTATGATTATTTGATG  
TATGCCGACATCGATTATGACCATCCTGA TGTCGCAGCAGAAATTAAGAGATGGGG  
CACTTGGTATGCCAATGAACTGCAATTGGACGGTTTCCGTCTTGATGCTGTCA  
AACACATTAAATTTTCTTTTTTTCGCGGATTGGGTTAATCATGTCAGGGAAAAAAC  
GGGGAAGGAAATGTTTACGGTAGCT GAATATTGGCAGAATGACTTGGGCGCGCT  
GGAAAAC TATTTGAACAAAACAAATTTTAATCATT CAGTGTTTGACGTGCC  
GCTTCATTATCAGTTCCATGCTGCATCGACACAGGGAGGCGGCTATGATATGAGG  
AAATTGCTGAACGGTACGGTCGTTTCCAAGCATCCGTTGAAATCGGTTACATTTGT

CGATAACCATGATACACAGCCGGGGCAATCGCTTGAGTCGACTGTCCAA  
ACATGGTTTAAGCCGCTTGCTTACGCTTTTATTCTCACAAGGGAATCTGGATACC  
CTCAGGTTTTCTACGGGGATATGTACGGGACGAAAGGAGACTCCCAGCGCGAAAT  
TCCTGCCTTGAAACACAAAATTGAACCGATCTTAAAAGCGAGAAAACAGT  
5 ATGCGTACGGAGCACAGCATGATTATTTGACCACCATGACATTGTCGGCTGGAC  
AAGGGAAGGCGACAGCTCGGTTGCA AATTCAGGTTTGGCGGCATTAATAACAGA  
CGGACCCGGTGGGGCAAAGCGAATGTATGTCGGCCGGCAAACGCCGGTGA  
GACATGGCATGACATTACCGGAAACCGTTCGGAGCCGGTTGTCATCAATTCCGA  
AGGCTGGGGAGAGTTTCACGTAAACG GCGGGTCGGTTTCAATTTATGTTCAAAGAT  
10 AGaagagcagagaggacggatttcctgaaggaaatccgttttttattt

SEQ ID NO: 2

ANLNGTLMQYFEWYMPNDGQHWRLQNDSAYLAEHGITAV  
WIPPAYKGTSQLADVGYGAYDLYDLGEFHQKGTVRTKYGTK  
5 GELQSAIKSLHSRDINVYGDVVINHKGGADATEDVTAVEV  
DPADRNRVISGEHLIKAWTHFHFPGRGSTYSDFKWHWYHF  
DGTDWDESRLNRIYKFQGKAWDWEVSNENGN DYLMYAD  
IDYDHPDVAAEIKRWGTWYANELQLDGFRLDAVKHIKFSF  
LRDWVNVHREKTGKEMFTVAEYWQNDLGALENYLNKTNFN  
20 HSVFDVPLHYQFHAASQTGGGYDMRKLLNGTVVSKHPLKS  
VTFVDNHDTQPGQSLESTVQTWFKPLAYAFILTRESGY PQ  
VFYGD MYG TKGDSQREIPALKHKIEPILKARKQYAYGAQH  
DYFDHHDIVGWTREGDSSVANSGLAALITDGP GGAKRMYV  
GRQNA GETWHDITGNRSEPVVINSEGWGEFHVNGGSVSIY  
25 VQR

SEQ ID NO: 3

gccccgcacatacgaagactggctgaaaacattgagcctttgatgactgatgattggctgaagaagtggatcgattg  
tttgagaaaagaagaagaccataaaaaatccttgtctgtcatcagacagggtatttttatgctgtccagactgtccgct  
30 gtgtaaaaataaggaataaaggggggtgttatttttactgatatgtaaaatataattgtataagaaaatgagagggg agaggaaac

0937563.06099  
66099" 994360

atgattcaaaaacgaaagcggacagtttcgttcagacttgtgcttatgtgcacgctgttatttgcagttt  
gccgattacaaaaacatcagccGTAAATGGCACGCTGATGCAGTATTTTGAATGGTATACGCC  
GAACGACGGCCAGCATGGAAACGATTGCAGAATGATGCGGAACATTTATCGGAT  
ATCGGAATCACTGCCGTCTGGATTCTCCCGCATACAAAGGA TTGAGCCAATCCGA  
5 TAACGGATACGGACCTTATGATTTGTATGATTTAGGAGAATTCCAGCAAAAAGG  
GACGGTCAGAACGAAATACGGCACAAAATCAGAGCTTCAAGATGCGATCGGCTCAC  
TGCATTCCCGGAACGTCCAAGTATACGGAGATGTGG TTTTGAATCATAAGGCTGGT  
GCTGATGCAACAGAAGATGTAAGTCCGTCGAAGTCAATCCGGCCAATAGAAAT  
CAGGAA ACTTCGGAGGAATATCAAATCAAAGCGTGGACGGATTTTCGTTTTCCGGG  
10 CCGTGGAAACACGTACAGTGATTTTAAATG GCATTGGTATCATTTTCGACGGAGCGG  
ACTGGGATGAATCCCGGAAGATCAGCCGCATCTTTAAGTTTCGTGGGGAAGGAA  
AAGCGTGGGATTGGGAAGTATCAAGTGAAAACGGCAACTATGACTATTTAATGT  
ATGCTGATGTTGACTACGACCACCTGATGTCTGTGGCAGAGACAAAAAATGGGG  
TATCTGGTATGCGAATGAACTGTCATTAGACGGCTTCCGTATTGATGCCGC  
CAAACATATTAAATTTTCATTTCTGCGTGATTGGGTTCAGGCGGTCAGACAGGCG  
ACGGGAAAAGAAATGTTTACGGTTG CGGAGTATTGGCAGAATAATGCCGGGAAA  
CTCGAAAACACTACTTGAATAAAACAAGCTTTAATCAATCCGTGTTTGATGTT  
CCGCTTCATTTCAATTTACAGGCGGCTTCCTCACAAGGAGGCGGATATGATATGA  
GGCGTTTGCTGGACGGTACCGTTGTGTCCAGGCATCCGGAAAAGGCGGTTACATT  
20 TGTTGAAAATCATGACACACAGCCGGGACAGTCATTGGAATCGACAGTCC  
AACTTGGTTTAAACCGCTTGCATACGCCTTTATTTTGACAAGAGAATCCGGTTA  
TCCTCAGGTGTTCTATGGGGATATG TACGGGACAAAAGGGACATCGCCAAAGGA  
AATTCCCTCACTGAAAGATAATATAGAGCCGATTTTAAAAGCGCGTAAGGA  
GTACGCATACGGGCCCCAGCACGATTATATTGACCACCCGGATGTGATCGGATGG  
25 ACGAGGGAAGGTGACAGCTCCGCCGCCAAATCAGGTTTGGCCGCTTTAATCACGG  
ACGGACCCGGCGGATCAAAGCGGATGTATGCCGGCCTGAAAAATGCCGGC  
GAGACATGGTATGACATAACGGGCAACCGTTCAGATACTGTAAAAATCGGATCT  
GACGGCTGGGGAGAGTTTCATGTAAA CGATGGGTCCGTCTCCATTTATGTTCAGA  
AATAA  
30 g g t a a t a a a a a a c a c c t c c a a g c t g a g t g c g g g t a t c a g c t t g g a  
gggtgcgtttatttttcagccgtatgacaaggtcggcatcaggtgtgacaaatacggatgctggctgtcataggtgaca

aatccggggttttgcgccgtttggctttttcacatgtctgattttgtataatcaacaggcacggagccggaatctttcgc  
cttgaaaaataagcggcgatcgtagctgcttccaatatggattgttcacgggatcgctgcttttaacacaacgtggg atcc

SEQ ID NO: 4

5 VNGTLMQYFEWYTPNDGQHWKRLQNDAEHLSDIGITAVWI  
PPAYKGLSQSDNGYGPYDL YDLGEFQQKGTVRTKYGTKSE  
LQDAIGSLHSRNVQVYGDVVLNHNKAGADATEDVTAVEVNP  
ANRNQETSEEYQIKAWTDFRFPGRGNTYSDFKWHWYHFDG  
ADWDESRKISRIFKFRGEGKAWDWEVSSSENGNYDYL MYAD  
10 VDYDHPDVVAETKKWGIWYANELSLDGFRIDA AKHIKFSF  
LRDWVQAVRQATGKEMFTVAEYWQNNAGKLENYLNKTSFN  
QSVFDVPLHFNLQAASSQGGGYDMRRLLDGTVVSRHPEKA  
VTFVENHDTQPGQSLESTVQTFWKPLAYAFILTRESGY PQ  
VFYGDMYGTGKTSPKEIPSLKDNIEPILKARKEYAYGPQH  
15 DYIDHPDVIGWTREGDSSAAKSGLAALITDGP GGSKRMYA  
GLKNAGETWYDITGNRSDTVKIGSDGWGEFHVNDGSVSIY

SEQ ID NO: 5

20 aaattcgatattgaaaacgattacaaataaaaaattataatagacgtaaacgttcgaggggttgctcccttttactcttt  
ttatgcaatcgtttcccttaatttttgaagccaaaccgtcgaatgtaacatttgattaagggggaagggcatt  
gtgct aacgtttcacgcacatcattcgaaaaggatggatgttcctgctcgcgttttgctcactgtctcgctgttctgcccacag  
gacagcccgccaaggctGCCGCACCGTTTAACGGCACCATGATGCAGTATTTTGAATGGTA  
CTTGCCGGATGATGGCACG TTATGGACCAAAGTGGCCAATGAAGCCAACA ACTTAT  
CCAGCCTTGGCATCACCGCTCTTTGGCTGCCGCCCGCTTACAA  
25 AGGAACAAGCCGCAGCGACGTAGGGTACGGAGTATACGACTTGTATGACCTCGG  
CGAATTCAATCAAAAAGGGACCGTCC GCACAAAATACGGAACAAAAGCTCAATA  
TCTTCAAGCCATTCAAGCCGCCACGCGCTGGAATGCAAGTGTACGCCGAT  
GTCGTGTTTCGACCATAAAGGCGGCGCTGACGGCACGGAATGGGTGGACGCCGTC  
GAAGTCAATCCGTCCGACCGCAACCAAGAAATCTCGGGCACCTATCAAATCCAAG  
30 CATGGACGAAATTTGATTTTCCCGGGCGGGGCAACACCTACTCCAGCTTTA  
AGTGGCGCTGGTACCATTTTGACGGCGTTGATTGGGACGAAAGCCGAAAATTGA  
GCCGCATTTACAAATTCCGCGGCATC GGCAAAGCGTGGGATTGGGAAGTAGAC

ACGGAAAACGGAAACTATGACTACTTAATGTATGCCGACCTTGATATGGATCA  
 TCCCGAAGTCGTGACCGAGCTGAAAAACTGGGGGAAATGGTATGTCAACACAAC  
 GAACATTGATGGGTTCCGGCTTGATGCCGTCAAGCATATTAAGTTCAGTTTTTTC  
 CTGATTGGTTGTCGTATGTGCGTTCTCAGACTGGCAAGCCGCTATTTACC  
 5 GTCGGGGAATATTGGAGCTATGACATCAACAAGTTGCACAATTACATTACGAAA  
 ACAGACGGAACGATGTCTTTGTTTGATGCCCCGTTACACAACAAATTTTATACCGC  
 TTCCAAATCAGGGGGCGCATTGATATGCGCACGTTAATGACCAATACTC  
 TCATGAAAGATCAACCGACATTGGCCGTCACCTTCGTTGATAATCATGACACCGA  
 ACCCGGCCAAGCGCTGCAGTCATGG GTCGACCCATGGTTCAAACCGTTGGCTTACG  
 10 CCTTTATTCTAACTCGGCAGGAAGGATACCCGTGCGTCTTTTATGGTGA  
 CTATTATGGCATTCCACAATATAACATTCCTTCGCTGAAAAGCAAATCGATCCG  
 CTCCTCATCGCGCGCAGGGATTATGTTACGGAACGCAACATGATTATCTTGATCA  
 CTCCGACATCATCGGGTGGACAAGGGAAGGGGGGCACTGAAAAACCAGGA  
 TCCGGACTGGCCGCACTGATACCGATGGGCCGGGAGGAAGCAAATGGATGTAC  
 GTTGGCAAACAACACGCTGGAAAAGT GTTCTATGACCTTACCGGCAACCGGAGT  
 GACACCGTCACCATCAACAGTGATGGATGGGGGGGAATTCAAAGTCAATGGCG  
 GTTCGGTTTCGGTTTGGGTTCTAGAAAAACGACCGTTTCTACCATCGCTCGGCC  
 GATCACAACCCGACCGTGGACTGGTGAATTCGTCCGTTGGACCGAACCACGGTTG  
 GTGGCATGGCCTTGA

tcctgcga

SEQ ID NO: 6

AAPFNGTMMQYFEWYLPDDGTLWTKVANEANNLSSLGITA  
 LWLPPAYKGTSRSDVGYGVYDLIDLGEFNQKGTVRTKYGT  
 25 KAQYLQAIQAAHAAGMQVYADVVDHKGGADGTEWVDAVE  
 VNPSDRNQEISGTYQIQAWTKFDFPGRGNTYSSFKWRWYH  
 FDGVDWDESRKLSRIYKFRGIGKAWDWEVDTENGNIDYLM  
 YADLMDMHPEVVTELKNWGKWWYVNTTNIDGFRLDAVKHIK  
 FSFFPDWLSYVRSQTGKPLFTVGEYWSYDINKLHNYITKT  
 30 DGTMSLFDAPLHNKFYTASKSGGAFDMRTLMTNTLMKDQP  
 TLAVTFVDNHDTEPGQALQSWVDPWFKPLAYAFILTRQEG  
 YPCVFYGDYYGIPQYNIPSLKSKIDPLLIARRDYAYGTQH

DYLDHSDIIGWTREGGTEKPGSGLAALITDGPGGSKWMYV  
GKQHAGKVFYDLTGNRSDTVTINS DGWGEFKVNGGSVSVW  
VPRKTTVSTIARPITTRPWTGEFVRWTEPRLVAW

5 SEQ ID NO: 10

1 ATPADWRSQS IYFLLTDRFA RTDGSTTATC  
31 NTADQKYCGG TWQGHIKLD YIQGMGFTAI  
61 WITPVTALP QTTAYGDAYH GYWQQDIYSL  
91 NENYGTADDL KALSSALHER GMYLMVDVVA  
121 NHMGYDGAGS SVDYSVFKPF SSQDYFHPFC  
151 FIQNYEDQTQ VEDCWLGDNT VSLPDLDTTK  
181 DVVKNEWYDW VGSLVSNYSI DGLRIDTVKH  
211 VQKDFWPGYN KAAGVYCIGE VLDGDPAYTC  
241 PYQNVMDGVL NYPIYYPLLN AFKSTSGSMD  
271 DLYNMINTVK SDCPDSTLLG TFVENHDNPR  
301 FASYTNDIAL AKNVAAFIIL NDGIPIYAG  
331 QEQHYAGGND PANREATWLS GYPTDSELYK  
361 LIASANAIRN YAISKDTGFV TYKNWPIYKD  
391 DITIAMRKGT DGSQIVTILS NKGASGDSYT  
421 LSLSGAGYTA GQQLTEVIGC TTVTVGSDGN  
451 VPVPMAGGLP RVLYPTEKLA GSKICSSS

668990" 6952450



## CLAIMS

1. A method of constructing a variant of a parent Termamyl-like  $\alpha$ -amylase, which variant has  $\alpha$ -amylase activity and at least one altered property as compared to said parent  $\alpha$ -amylase, which method comprises

i) analysing the structure of the parent Termamyl-like  $\alpha$ -amylase to identify at least one amino acid residue or at least one structural part of the Termamyl-like  $\alpha$ -amylase structure, which amino acid residue or structural part is believed to be of relevance for altering said property of the parent Termamyl-like  $\alpha$ -amylase (as evaluated on the basis of structural or functional considerations),

ii) constructing a Termamyl-like  $\alpha$ -amylase variant, which as compared to the parent Termamyl-like  $\alpha$ -amylase, has been modified in the amino acid residue or structural part identified in i) so as to alter said property; and, optionally,

iii) testing the resulting Termamyl-like  $\alpha$ -amylase variant with respect to said property or properties.

2. The method according to claim 1, wherein the property to be altered is selected from the group consisting of substrate specificity, substrate binding, substrate cleavage pattern, temperature stability, pH dependent activity, pH dependent stability (especially increased stability at low (e.g. pH<6) or high (e.g. pH>9) pH values), stability towards oxidation, Ca<sup>2+</sup>-dependency and specific activity.

3. The method according to claim 1, wherein the property to be altered is the calcium ion dependency and the structural part to be modified is selected from the group consisting of the C domain, the interface between the A and B domain, the interface between the A and C domain, or the interaction to a calcium binding site of the Termamyl-like  $\alpha$ -amylase.

4. The method according to claim 1, wherein the property to be altered is the substrate cleavage pattern and the structural part to be modified is located within 10Å from an amino acid residue of the substrate binding site.

5. A method of constructing a variant of a parent Termamyl-like  $\alpha$ -amylase, which variant has  $\alpha$ -amylase activity and one or more altered properties as compared to said parent  $\alpha$ -amylase, which method comprises

i) comparing the three-dimensional structure of the Termamyl-like  $\alpha$ -amylase with the structure of a non-Termamyl-like  $\alpha$ -amylase,

ii) identifying a part of the Termamyl-like  $\alpha$ -amylase structure which is different from the non-Termamyl-like  $\alpha$ -amylase structure and which from structural or functional considerations is contemplated to be responsible for differences in one or more properties of the Termamyl-like and non-Termamyl-like  $\alpha$ -amylase,

iii) modifying the part of the Termamyl-like  $\alpha$ -amylase identified in ii) whereby a Termamyl-like  $\alpha$ -amylase variant is obtained, one or more properties of which differ from the parent Termamyl-like  $\alpha$ -amylase, and optionally,

iv) testing the resulting Termamyl-like  $\alpha$ -amylase variant with respect to said property or properties.

6. The method according to claim 5, wherein, in step iii), the part of the Termamyl-like  $\alpha$ -amylase is modified so as to resemble the corresponding part of the non-Termamyl-like  $\alpha$ -amylase.

7. The method according to claim 5, wherein, in step iii), the modification is accomplished by deleting one or more amino acid residues of the part of the Termamyl-like  $\alpha$ -amylase to be modified; by replacing one or more amino acid residues of the part of the Termamyl-like  $\alpha$ -amylase to be modified with the amino acid residues occupying corresponding positions in the non-Termamyl-like  $\alpha$ -amylase; or by insertion of one or more amino acid residues present in the non-Termamyl-like  $\alpha$ -amylase into a corresponding position in the Termamyl-like  $\alpha$ -amylase.

8. The method according to claim 5, wherein the non-Termamyl-like  $\alpha$ -amylase structure is the structure of a fungal  $\alpha$ -amylase or a mammalian  $\alpha$ -amylase.

9. The method according to claim 8, wherein the non-Termamyl-like  $\alpha$ -amylase is the *Aspergillus oryzae* TAKA  $\alpha$ -amylase, the *A. niger* acid  $\alpha$ -amylase, the *Bacillus subtilis*  $\alpha$ -amylase or the pig pancreatic  $\alpha$ -amylase.

10. The method according to claim 1, wherein the parent Termamyl-like  $\alpha$ -amylase is derived from a strain of *Bacillus*.

11. The method according to claim 10, wherein the parent  $\alpha$ -amylase is derived from a strain of a *B. licheniformis*, *B. amyloliquefaciens*, *B. stearothermophilus* or a strain from an alkalophilic *Bacillus* sp. such as NCIB 12289, NCIB 12512 or NCIB 12513.

12. The method according to claim 1, wherein the parent  $\alpha$ -amylase is a hybrid  $\alpha$ -amylase comprising a combination of partial amino acid sequences derived from at least two  $\alpha$ -amylases, of which one is a Termamyl-like  $\alpha$ -amylase and the other(s) are, e.g., from a microbial and/or a mammalian  $\alpha$ -amylase.

13. The method according to claim 5, wherein the part of the parent Termamyl-like  $\alpha$ -amylase to be modified and identified in step ii) is loop 1, loop 2, loop 3 and/or loop 8 of the parent  $\alpha$ -amylase.

13. A method of constructing a variant of a parent Termamyl-like  $\alpha$ -amylase, which has a decreased calcium ion dependency as compared to said parent, which method comprises:

i) identifying an amino acid residue within 10Å from a  $\text{Ca}^{2+}$  binding site of a Termamyl-like  $\alpha$ -amylase in a model of the three-dimensional structure of said  $\alpha$ -amylase, which from structural or functional considerations is believed to be responsible for a non-optimal calcium ion interaction,

ii) constructing a variant in which said amino acid residue is replaced with another amino acid residue which from structural or functional considerations is believed to be important for establishing a higher  $\text{Ca}^{2+}$  binding affinity, and

iii) testing the  $\text{Ca}^{2+}$  dependency of the resulting Termamyl-like  $\alpha$ -amylase variant.

14. A method of constructing a variant of a parent Termamyl-like  $\alpha$ -amylase which variant has  $\alpha$ -amylase activity and an altered pH dependent activity, which method comprises

i) in a three-dimensional structure of the Termamyl-like  $\alpha$ -amylase in question, identifying an amino acid residue within 15Å from an active site residue, in particular 10Å from an active site residue, which amino acid residue is contemplated to be involved in electrostatic or hydrophobic interactions with an active site residue,

ii) replacing, in the structure, said amino acid residue with an amino acid residue which changes the electrostatic and/or hydrophobic surroundings of an active site residue and evaluating the accomodation of the amino acid residue in the structure,

iii) optionally repeating step i) and/or ii) until an amino acid replacement has been identified which is accomodated into the structure,

iv) constructing a Termamyl-like  $\alpha$ -amylase variant resulting from steps i), ii) and optionally iii) and testing the pH dependent activity of said variant.

15. A method of increasing the thermostability and/or altering the temperature optimum of a parent Termamyl-like  $\alpha$ -amylase, which method comprises

i) identifying an internal hole or a crevice of the parent Termamyl-like  $\alpha$ -amylase in the three-dimensional structure of said  $\alpha$ -amylase,

ii) replacing, in the structure, one or more amino acid residues in the neighbourhood of the hole or crevice identified in i) with another amino acid residue which from structural or functional considerations is believed to increase the hydrophobic interaction and to fill out or reduce the size of the hole or crevice,

iii) constructing a Termamyl-like  $\alpha$ -amylase variant resulting from step ii) and testing the thermostability and/or temperature optimum of the variant.

16. A method of constructing a variant of a Termamyl-like  $\alpha$ -amylase which has a reduced ability to cleave a substrate close to the branching point, which method comprises

i) identifying the substrate binding area of the parent Termamyl-like  $\alpha$ -amylase in a model of the three-dimensional structure of said  $\alpha$ -amylase,

ii) replacing, in the model, one or more amino acid residues of the substrate binding area of the cleft identified in i), which is/are believed to be responsible for the cleavage pattern of the parent  $\alpha$ -amylase, with another amino acid residue which from

structural considerations is believed to result in an altered substrate cleavage pattern, or deleting one or more amino acid residues of the substrate binding area contemplated to introduce favourable interactions to the substrate or adding one or more amino acid residues to the substrate binding area contemplated to introduce favourable interactions to the substrate, and

iii) constructing a Termamyl-like  $\alpha$ -amylase variant resulting from step ii) and testing the substrate cleavage pattern of the variant.

17. The method according to claim 1, in which the  $\alpha$ -amylase variant is obtained by cultivating a microorganism comprising a DNA sequence encoding the variant under conditions which are conducive for producing the variant, and optionally subsequently recovering the variant from the resulting culture broth.

18. A variant of a parent Termamyl-like  $\alpha$ -amylase, in which variant at least one amino acid residue of the parent  $\alpha$ -amylase, which is/are present in a fragment corresponding to the amino acid fragment 44-57 of the amino acid sequence of SEQ ID NO: 4, has been deleted or replaced with one or more amino acid residues which is/are present in a fragment corresponding to the amino acid fragment 66-84 of the amino acid sequence shown in SEQ ID NO: 10, or in which one or more additional amino acid residues has been added using the relevant part of SEQ ID NO: 10 or a corresponding part of another Fungamyl-like  $\alpha$ -amylase as a template.

19. A variant of a parent Termamyl-like  $\alpha$ -amylase, which variant has a region which, when the amino acid sequence of variant is aligned most closely with the amino acid sequence of the said parent  $\alpha$ -amylase, occupies the same position as the portion from residue X to residue Y of SEQ ID NO: 4, the said region having at least 80% sequence homology with the part of SEQ ID NO: 10 extending from residue Z to residue V of SEQ ID NO: 10, wherein

X is the amino acid residue occupying position 44, 45, 46, 47 or 48 of SEQ ID NO: 4,

Y is the amino acid residue occupying position 51, 52, 53, 54, 55, 56 or 57 of SEQ ID NO: 4,

Z is the amino acid residue occupying position 66, 67, 68, 69 or 70 of SEQ ID NO: 10, and

V is the amino acid residue occupying position 78, 79, 80, 81, 82, 83 or 84 of SEQ ID NO: 10.

20. The variant according to claim 18, wherein X is the amino acid residue occupying position 48 and Y the amino acid residue occupying position 51 of SEQ ID NO: 4 and Z is the amino acid residue occupying position 70 and V the amino acid residue occupying position 78 in SEQ ID NO: 10.

21. A variant of a parent Termamyl-like  $\alpha$ -amylase, in which variant at least one of the amino acid residues of the parent  $\alpha$ -amylase, which is/are present in an amino acid fragment corresponding to the amino acid fragment 195-202 of the amino acid sequence of SEQ ID NO: 4, has been deleted or replaced with one or more of the amino acid residues which is/are present in an amino acid fragment corresponding to the amino acid fragment 165-177 of the amino acid sequence shown in SEQ ID NO: 10, or in which one or more additional amino acid residues has been added using the relevant part of SEQ ID NO: 10 or a corresponding part of another Fungamyl-like  $\alpha$ -amylase as a template.

22. A variant of a parent Termamyl-like  $\alpha$ -amylase, which variant has a region which, when the amino acid sequence of variant is aligned most closely with the amino acid sequence of the said parent  $\alpha$ -amylase, occupies the same position as the portion from residue X to residue Y of SEQ ID NO: 4, the said region having at least 80%, such as 90% sequence homology with the part of SEQ ID NO: 10 extending from residue Z to residue V of SEQ ID NO: 10, wherein

X is the amino acid occupying position 195 or 196 of SEQ ID NO: 4,

Y is the amino acid residue occupying position 198, 199, 200, 201, or 202 of SEQ ID NO: 4,

Z is the amino acid residue occupying position 165 or 166 of SEQ ID NO: 10,

and

V is the amino acid residue occupying position 173, 174, 175, 176 or 177 of SEQ ID NO: 10.

23. The variant according to claim 21, in which the amino acid fragment of the parent  $\alpha$ -amylase, which corresponds to amino acid residues 196-198 of SEQ ID NO: 4, has been replaced with the amino acid fragment corresponding to amino acid residues 166-173 of the amino acid sequence shown in SEQ ID NO: 10.

24. A variant of a parent Termamyl-like  $\alpha$ -amylase, in which variant at least one of the amino acid residues of the parent  $\alpha$ -amylase, which is/are present in a fragment corresponding to the amino acid fragment 117-185 of the amino acid sequence of SEQ ID NO: 4, has/have been deleted or replaced with one or more of the amino acid residues, which is/are present in an amino acid fragment corresponding to the amino acid fragment 98-210 of the amino acid sequence shown in SEQ ID NO: 10, or in which one or more additional amino acid residues has been added using the relevant part of SEQ ID NO: 10 or a corresponding part of another Fungamyl-like  $\alpha$ -amylase as a template.

25. A variant of a parent Termamyl-like  $\alpha$ -amylase, which variant has a region which, when the amino acid sequence of variant is aligned most closely with the amino acid sequence of the said parent  $\alpha$ -amylase, occupies the same position as the portion from residue X to residue Y of SEQ ID NO: 4, the said region having at least 80%, such as at least 90% sequence homology with the part of SEQ ID NO: 10 extending from residue Z to residue V of SEQ ID NO: 10, wherein

X is the amino acid occupying position 117, 118, 119, 120 or 121 of SEQ ID NO: 4,

Y is the amino acid occupying position 181, 182, 183, 184 or 185 of SEQ ID NO: 4,

Z is the amino acid occupying position 98, 99, 100, 101, 102 of SEQ ID NO: 10, and

V is the amino acid occupying position 206, 207, 208, 209 or 210 of SEQ ID NO: 10.

26. The variant according to claim 24, in which an amino acid fragment of the parent  $\alpha$ -amylase, which corresponds to amino acid residues 121-181 of SEQ ID NO: 4, has been

replaced with the amino acid fragment corresponding to amino acid residues 102-206 of the amino acid sequence shown in SEQ ID NO: 10.

27. A variant of a parent Termamyl-like  $\alpha$ -amylase, in which variant at least one of the amino acid residues of the parent  $\alpha$ -amylase, which is/are present in a fragment corresponding to the amino acid fragment 117-181 of the amino acid sequence of SEQ ID NO: 4, has/have been deleted or replaced with one or more of the amino acid residues, which is/are present in an amino acid fragment corresponding to the amino acid fragment to 98-206 of the amino acid sequence shown in SEQ ID NO: 10, or in which one or more additional amino acid residues has been added using the relevant part of SEQ ID NO: 10 or a corresponding part of another Fungamyl-like  $\alpha$ -amylase as a template.

28. A variant of a parent Termamyl-like  $\alpha$ -amylase, which variant has a region which, when the amino acid sequence of variant is aligned most closely with the amino acid sequence of the said parent  $\alpha$ -amylase, occupies the same position as the portion from residue X to residue Y of SEQ ID NO: 4, the said region having at least 80%, such as at least 90% sequence homology with the part of SEQ ID NO: 10 extending from residue Z to residue V of SEQ ID NO: 10, wherein

X is the amino acid occupying position 117, 118, 119, 120 or 121 of SEQ ID NO: 4,

Y is the amino acid occupying position 174, 175, 176 or 177 of SEQ ID NO: 4,

Z is the amino acid occupying position 98, 99, 100, 101, 102 of SEQ ID NO: 10, and

V is the amino acid occupying position 199, 200, 201 or 202 of SEQ ID NO: 10.

29. The variant according to claim 27, in which the amino acid fragment of the parent  $\alpha$ -amylase, which corresponds to amino acid residues 121-174 of SEQ ID NO: 4, has been replaced with the amino acid fragment corresponding to amino acid residues 102-199 of the amino acid sequence shown in SEQ ID NO: 10.



30. A variant of a parent Termamyl-like  $\alpha$ -amylase, in which variant at least one of the amino acid residues of the parent  $\alpha$ -amylase, which is/are present in an amino acid fragment corresponding to the amino acid fragment 12-19 of the amino acid sequence of SEQ ID NO: 4, has/have been deleted or replaced with one or more of the amino acid residues, which is/are present in an amino acid fragment which corresponds to the amino acid fragment 28-42 of SEQ ID NO: 10, or in which one or more additional amino acid residues has/have been inserted using the relevant part of SEQ ID NO: 10 or a corresponding part of another Fungamyl-like  $\alpha$ -amylase as a template.

31. A variant of a parent Termamyl-like  $\alpha$ -amylase, which variant has a region which, when the amino acid sequence of variant is aligned most closely with the amino acid sequence of the said parent  $\alpha$ -amylase, occupies the same position as the portion from residue X to residue Y of SEQ ID NO: 4, the said region having at least 80%, such as at least 90% sequence homology with the part of SEQ ID NO: 10 extending from residue Z to residue V of SEQ ID NO: 10, wherein

X is the amino acid occupying position 12, 13 or 14 of SEQ ID NO: 4,

Y is the amino acid occupying position 15, 16, 17, 18 or 19 of SEQ ID NO:

4,

Z is the amino acid occupying position 28, 29, 30, 31 or 32 of SEQ ID NO:

10, and

V is an amino acid residue corresponding to the amino acid occupying position 38, 39, 40, 41 or 42 of SEQ ID NO: 10.

32. The variant according to claim 30, in which the amino acid fragment of the parent  $\alpha$ -amylase, which corresponds to amino acid residues 14-15 of SEQ ID NO: 4, has been replaced with the amino acid fragment corresponding to amino acid residues 32-38 of the amino acid sequence shown in SEQ ID NO: 10.

33. A variant of a parent Termamyl-like  $\alpha$ -amylase, in which variant at least one of the amino acid residues of the parent  $\alpha$ -amylase, which is present in a fragment corresponding to amino acid residues 7-23 of the amino acid sequence of SEQ ID NO: 4, has/have been deleted or replaced with one or more amino acid residues, which is/are present in an amino

acid fragment corresponding to amino acid residues 13-45 of the amino acid sequence shown in SEQ ID NO: 10, or or in which one or more additional amino acid residues has/have been inserted using the relevant part of SEQ ID NO: 10 or a corresponding part of another Fungamyl-like  $\alpha$ -amylase as a template.

5

34. A variant of a parent Termamyl-like  $\alpha$ -amylase, which variant has a region which, when the amino acid sequence of variant is aligned most closely with the amino acid sequence of the said parent  $\alpha$ -amylase, occupies the same position as the portion from residue X to residue Y of SEQ ID NO: 4, the said region having at least 80%, such as at least 90% sequence  
10 homology with the part of SEQ ID NO: 10 extending from residue Z to residue V of SEQ ID NO: 10, wherein

X is the amino acid occupying position 7 or 8 of SEQ ID NO: 4,

Y is the amino acid occupying position 18, 19, 20, 21, 22 or 23 of SEQ ID  
NO: 4,

Z is the amino acid occupying position 13 or 14 of SEQ ID NO: 10, and

V is the amino acid occupying position 40, 41, 42, 43, 44 or 45 of SEQ ID  
NO: 10.

35. The variant according to claim 33, in which the amino acid fragment of the parent  $\alpha$ -amylase, which corresponds to amino acid residues 8-18 of SEQ ID NO: 4, has been replaced,  
20 with the amino acid fragment corresponding to amino acid residues 14-40 of the amino acid sequence shown in SEQ ID NO: 10.

36. A variant of a parent Termamyl-like  $\alpha$ -amylase, in which variant at least one of the  
25 amino acid residues of the parent  $\alpha$ -amylase, which is present in a fragment corresponding to amino acid residues 322-346 of the amino acid sequence of SEQ ID NO: 2, has/have been deleted or replaced with one or more amino acid residues, which is/are present in an amino acid fragment corresponding to amino acid residues 291-313 of the amino acid sequence shown in SEQ ID NO: 10, or or in which one or more additional amino acid residues  
30 has/have been inserted using the relevant part of SEQ ID NO: 10 or a corresponding part of another Fungamyl-like  $\alpha$ -amylase as a template.

37. A variant of a parent Termamyl-like  $\alpha$ -amylase, which variant has a region which, when the amino acid sequence of variant is aligned most closely with the amino acid sequence of the said parent  $\alpha$ -amylase, occupies the same position as the portion from residue X to residue Y of SEQ ID NO: 2, the said region having at least 80% sequence homology with the part of SEQ ID NO: 10 extending from residue Z to residue V of SEQ ID NO: 10, wherein

X is the amino acid occupying position 322, 323, 324 or 325 of SEQ ID NO:

2,

Y is the amino acid occupying position 343, 344, 345 or 346 of SEQ ID NO:

2,

Z is the amino acid occupying position 291, 292, 293 or 294 of SEQ ID NO:

10, and

V is the amino acid occupying position 310, 311, 312 or 313 of SEQ ID NO:

10.

38. The variant according to claim 36, in which the amino acid fragment of the parent  $\alpha$ -amylase, which corresponds to amino acid residues 325-345 of SEQ D No. 2, has been replaced with the amino acid fragment corresponding to amino acid residues 294-313 of the amino acid sequence shown in SEQ ID NO: 10.

39. A variant of a parent Fungamyl-like  $\alpha$ -amylase, in which variant at least one of the amino acid residues of the parent  $\alpha$ -amylase, which is/are present in an amino acid fragment corresponding to amino acid residues 291-313 of the amino acid sequence of SEQ ID NO: 10, has/have been deleted or replaced with one or more of the amino acid residues, which is/are present in an amino acid fragment corresponding to amino acid residues 98-210 of the amino acid sequence shown in SEQ ID NO: 4, or in which one or more additional amino acid residues has/have been inserted using the relevant part of SEQ ID NO: 4 or a corresponding part of another Termamyl-like  $\alpha$ -amylase as a template.

40. A variant of a parent Fungamyl-like  $\alpha$ -amylase, which variant has a region which, when the amino acid sequence of variant is aligned most closely with the amino acid sequence of the said parent  $\alpha$ -amylase, occupies the same position as the portion from residue X to residue Y of SEQ ID NO: 10, the said region having at least 80%, such as at least 90% sequence

homology with the part of SEQ ID NO: 10 extending from residue Z to residue V of SEQ ID NO: 4, wherein

X is the amino acid occupying position 117, 118, 119, 120 or 121 of SEQ ID NO: 10,

Y is the amino acid occupying position 181, 182, 183, 184 or 185 of SEQ ID NO: 10,

Z is the amino acid occupying position 98, 99, 100, 101 or 102 of SEQ ID NO: 4, and

V is the amino acid occupying position 206, 207, 208, 209 or 210 of SEQ ID NO: 4.

41. The variant according to claim 39, in which the amino acid fragment of the parent  $\alpha$ -amylase, which corresponds to amino acid residues 121-181 of SEQ ID NO: 10, has been replaced with the amino acid fragment corresponding to amino acid residues 102-206 of the amino acid sequence shown in SEQ ID NO: 4.

42. A variant according to claim 39, in which the the amino acid fragment of the parent  $\alpha$ -amylase, which corresponds to amino acid residues 121-174 of SEQ ID NO: 10, has been replaced with the amino acid fragment corresponding to amino acid residues 102-199 of the amino acid sequence shown in SEQ ID NO: 4.

43. A variant of a parent Fungamyl-like  $\alpha$ -amylase, in which an amino acid fragment corresponding to amino acid residues 181-184 of the amino acid sequence shown in SEQ ID NO: 10 has been deleted.

44. A variant of a parent Termamyl-like  $\alpha$ -amylase, which exhibits  $\alpha$ -amylase activity and which has a decreased  $\text{Ca}^{2+}$  dependency as compared to the parent  $\alpha$ -amylase.

45. A variant according to claim 44, which comprises a mutation in a position corresponding to at least one of the following positions in SEQ ID NO 2:  
N104, A349, I479, L346, I430, N457, K385, F350, I411, H408 or G303, in particular a mutation corresponding to

N104D;  
A349C+I479C;  
L346C+I430C;  
N457D,E;  
5 N457D,E+K385R;  
F350D,E+I430R,K;  
F350D,E+I411R,K;  
H408Q,E,N,D; and/or  
G303N,D,Q,E.

10 46. A variant of a parent Termamyl-like  $\alpha$ -amylase which exhibits a higher activity below the pH optimum than the parent  $\alpha$ -amylase, which variant comprises a mutation of an amino acid residue corresponding to at least one of the following positions of the *B. licheniformis*  $\alpha$ -amylase (SEQ ID NO: 2): E336, Q333, P331, I236, V102, A232, I103, L196, in particular at least one of the following mutations:

E336R,K;  
Q333R,K; P331R,K;  
V102R,K,A,T,S,G;  
I236K,R,N;  
I103K,R;  
L196K,R; and/or  
A232T,S,G.

25 47. A variant of a parent Termamyl-like  $\alpha$ -amylase which exhibits a higher activity above the pH optimum than the parent  $\alpha$ -amylase, which variant comprises a mutation of an amino acid residue corresponding to at least one of the following positions of the *B. licheniformis*  $\alpha$ -amylase (SEQ ID NO: 2): N236, H281 and/or Y273, in particular one of the following mutations:

30 N326I,Y,F,L,V;  
H281F,I,L; and/or  
Y273F,W.

48. A variant of a parent Termamyl-like  $\alpha$ -amylase which exhibits  $\alpha$ -amylase activity and which has an increased thermostability and/or altered temperature optimum as compared to the parent  $\alpha$ -amylase, which variant comprises a mutation of an amino acid residue corresponding to at least one of the following positions of the *B. licheniformis*  $\alpha$ -amylase (SEQ ID NO: 2):

L61, Y62, F67, K106, G145, I212, S151, R214, Y150, F143, R146, L241, I236, L7, V259, F284, F350, F343, L427 and/or V481, in particular at least one of the following mutations:

L61W,V,F;

Y62W;

F67W;

K106R,F,W;

G145F,W

I212F,L,W,Y,R,K;

S151 replaced with any other amino acid residue and in particular with F,W,I or L;

R214W;

Y150R,K;

F143W;

R146W;

L241I,F,Y,W;

I236L,F,W,Y;

L7F,I,W;

V259F,I,L;

F284W;

F350W;

F343W;

L427F,L,W; and/or

V481,F,I,L,W.

49. A variant of a parent Termamyl-like  $\alpha$ -amylase, which exhibits  $\alpha$ -amylase activity and which has a reduced capability of cleaving an oligo-saccharide substrate close to the branching point as compared to the parent  $\alpha$ -amylase, which variant comprises a mutation of an amino

acid residue corresponding to at least one of the following positions of the *B. licheniformis*  $\alpha$ -amylase (SEQ ID NO: 2):

V54, D53, Y56, Q333 and/or G57, in particular at least one of the following mutations:

V54L,I,F,Y,W,R,K,H,E,Q;

D53L,I,F,Y,W;

Y,56W;

Q333W; and/or

G57 to all possible amino acid residues.

50. The variant according to claim 17, wherein one or more proline residues present in the amino acid residues with which the parent  $\alpha$ -amylase is modified are replaced with a non-proline residue such as alanine.

51. The variant according to claim 17, wherein one or more cysteine residues present in the amino acid residues with which the parent  $\alpha$ -amylase is modified are replaced with a non-cysteine residue such as alanine.

52. A DNA construct comprising a DNA sequence encoding an  $\alpha$ -amylase variant according to claim 17.

53. A recombinant expression vector which carries a DNA construct according to Claim 52.

54. A cell which is transformed with a DNA construct according to Claim 52.

55. A cell according to Claim 54, which is a microorganism.

56. A cell according to Claim 55, which is a bacterium or a fungus.

57. The cell according to Claim 56, which is a grampositive bacterium such as *Bacillus subtilis*, *Bacillus licheniformis*, *Bacillus lentus*, *Bacillus brevis*, *Bacillus stearothermophilus*, *Bacillus alkalophilus*, *Bacillus amyloliquefaciens*, *Bacillus coagulans*, *Bacillus circulans*, *Bacillus lautus* or *Bacillus thuringiensis*.

58. Use of an  $\alpha$ -amylase variant according to claim 17 for washing and/or dishwashing.

59. Use of an  $\alpha$ -amylase variant according to claim 17 for desizing.

5 60. Use of an  $\alpha$ -amylase variant according to claim 17 for starch liquefaction.

61. A detergent additive comprising an  $\alpha$ -amylase variant according to claim 17, optionally in the form of a non-dusting granulate, stabilised liquid or protected enzyme.

10 62. A detergent additive according to Claim 61 which contains 0.02-200 mg of enzyme protein/g of the additive.

63. A detergent additive according to Claim 61, which additionally comprises another enzyme such as a protease, a lipase, a peroxidase, another amylolytic enzyme and/or a cellulase.

64. A detergent composition comprising a surfactant and an  $\alpha$ -amylase variant according to claim 17.

65. A detergent composition according to Claim 64 which additionally comprises another enzyme such as a protease, a lipase, a peroxidase, another amylolytic enzyme and/or a cellulase.

66. A manual or automatic dishwashing detergent composition comprising an  $\alpha$ -amylase variant according to claim 17.

67. A dishwashing detergent composition according to Claim 66 which additionally comprises another enzyme such as a protease, a lipase, a peroxidase, another amylolytic enzyme and/or a cellulase.

68. A manual or automatic laundry washing composition comprising an  $\alpha$ -amylase variant according to claim 17.



69. A laundry washing composition according to Claim 68, which additionally comprises another enzyme such as a protease, a lipase, a peroxidase, an amylolytic enzyme and/or a cellulase.

## ABSTRACT

The present invention relates to a method of constructing a variant of a parent Termamyl-like  $\alpha$ -amylase, which variant has  $\alpha$ -amylase activity and at least one altered property as compared to the parent  $\alpha$ -amylase, comprises

i) analysing the structure of the parent Termamyl-like  $\alpha$ -amylase to identify at least one amino acid residue or at least one structural part of the Termamyl-like  $\alpha$ -amylase structure, which amino acid residue or structural part is believed to be of relevance for altering the property of the parent Termamyl-like  $\alpha$ -amylase (as evaluated on the basis of structural or functional considerations),

ii) constructing a Termamyl-like  $\alpha$ -amylase variant, which as compared to the parent Termamyl-like  $\alpha$ -amylase, has been modified in the amino acid residue or structural part identified in i) so as to alter the property, and, optionally,

iii) testing the resulting Termamyl-like  $\alpha$ -amylase variant with respect to the property in question.

1/11

66000" 0954260




Fig. 1

2/11

668090 6942660




Fig. 2

3/11

668090" 69542660




Fig. 3

4/11

668090" 6954260

LOOP1(a), TAA: 28-42, BA2:12-19




Fig. 4

5/11




Fig. 5

**Fig. 6**

1970-1971		1971-1972		1972-1973		1973-1974		1974-1975		1975-1976		1976-1977		1977-1978		1978-1979		1979-1980		1980-1981		1981-1982		1982-1983		1983-1984		1984-1985		1985-1986		1986-1987		1987-1988		1988-1989		1989-1990		1990-1991		1991-1992		1992-1993		1993-1994		1994-1995		1995-1996		1996-1997		1997-1998		1998-1999		1999-2000		2000-2001		2001-2002		2002-2003		2003-2004		2004-2005		2005-2006		2006-2007		2007-2008		2008-2009		2009-2010		2010-2011		2011-2012		2012-2013		2013-2014		2014-2015		2015-2016		2016-2017		2017-2018		2018-2019		2019-2020		2020-2021		2021-2022		2022-2023		2023-2024		2024-2025		2025-2026		2026-2027		2027-2028		2028-2029		2029-2030		2030-2031		2031-2032		2032-2033		2033-2034		2034-2035		2035-2036		2036-2037		2037-2038		2038-2039		2039-2040		2040-2041		2041-2042		2042-2043		2043-2044		2044-2045		2045-2046		2046-2047		2047-2048		2048-2049		2049-2050		2050-2051		2051-2052		2052-2053		2053-2054		2054-2055		2055-2056		2056-2057		2057-2058		2058-2059		2059-2060		2060-2061		2061-2062		2062-2063		2063-2064		2064-2065		2065-2066		2066-2067		2067-2068		2068-2069		2069-2070		2070-2071		2071-2072		2072-2073		2073-2074		2074-2075		2075-2076		2076-2077		2077-2078		2078-2079		2079-2080		2080-2081		2081-2082		2082-2083		2083-2084		2084-2085		2085-2086		2086-2087		2087-2088		2088-2089		2089-2090		2090-2091		2091-2092		2092-2093		2093-2094		2094-2095		2095-2096		2096-2097		2097-2098		2098-2099		2099-2100		2100-2101		2101-2102		2102-2103		2103-2104		2104-2105		2105-2106		2106-2107		2107-2108		2108-2109		2109-2110		2110-2111		2111-2112		2112-2113		2113-2114		2114-2115		2115-2116		2116-2117		2117-2118		2118-2119		2119-2120		2120-2121		2121-2122		2122-2123		2123-2124		2124-2125		2125-2126		2126-2127		2127-2128		2128-2129		2129-2130		2130-2131		2131-2132		2132-2133		2133-2134		2134-2135		2135-2136		2136-2137		2137-2138		2138-2139		2139-2140		2140-2141		2141-2142		2142-2143		2143-2144		2144-2145		2145-2146		2146-2147		2147-2148		2148-2149		2149-2150		2150-2151		2151-2152		2152-2153		2153-2154		2154-2155		2155-2156		2156-2157		2157-2158		2158-2159		2159-2160		2160-2161		2161-2162		2162-2163		2163-2164		2164-2165		2165-2166		2166-2167		2167-2168		2168-2169		2169-2170		2170-2171		2171-2172		2172-2173		2173-2174		2174-2175		2175-2176		2176-2177		2177-2178		2178-2179		2179-2180		2180-2181		2181-2182		2182-2183		2183-2184		2184-2185		2185-2186		2186-2187		2187-2188		2188-2189		2189-2190		2190-2191		2191-2192		2192-2193		2193-2194		2194-2195		2195-2196		2196-2197	
-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--



7/11




Fig. 7

66090" E954260

[illegible]

Fig. 8

9/11

301  
TCT AAT AGC GGT GGT TAT TAT GAT ATG AGA AAT ATT TTA AAT GGT TCT GTG GTG CAA AAA  
S N S G G Y Y D M R N I L N G S V V Q K

321  
CAT CCA ACA CAT GCC GTT ACT TTT GTT GAT AAC CAT GAT TCT CAG CCC GGG GAA GCA TTG  
H P T H A V T F V D N H D S Q P G E A L

341  
GAA TCC TTT GTT CAA CAA TGG TTT AAA CCA CTT GCA TAT GCA TTG GTT CTG ACA AGG GAA  
E S F V Q Q W F K P L A Y A L V L T R E

361  
CAA GGT TAT CCT TCC GTA TTT TAT GGG GAT TAC TAC GGT ATC CCA ACC CAT GGT GTT CCG  
Q G Y P S V F Y G D Y Y G I P T H G V P

381  
GCT ATG AAA TCT AAA ATA GAC CCT CTT CTG CAG GCA CGT CAA ACT TTT GCC TAT GGT ACG  
A M K S K I D P L L Q A R Q T F A Y G T

401  
CAG CAT GAT TAC TTT GAT CAT CAT GAT ATT ATC GGT TGG ACA AGA GAG GGA AAT AGC TCC  
Q H D Y F D H H D I I G W T R E G N S S

421  
CAT CCA AAT TCA GGC CTT GCC ACC ATT ATG TCA GAT GGT CCA GGT GGT AAC AAA TGG ATG  
H P N S G L A T I M S D G P G G N K W M

441  
TAT GTG GGG AAA AAT AAA GCG GGA CAA GTT TGG AGA GAT ATT ACC GGA AAT AGG ACA GGC  
Y V G K N K A G Q V W R D I T G N R T G

261  
ACC GTC ACA ATT AAT GCA GAC GGA TGG GGT AAT TTC TCT GTT AAT GGA GGG TCC GTT TCG  
T V T I N A D G W G N F S V N G G S V S

481  
GTT TGG GTG AAG CAA TAA  
V W V K Q

Fig. 8 (cont.)

10/11




Fig. 9

11/11




Fig. 10

As a below named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name.

I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled:

$\alpha$ -Amylase Mutants

the specification of which (check only one item below):

☐ is attached hereto

☒ was filed as United States application

Serial No. to be assigned

on July 18, 1996

and was amended

on \_\_\_\_\_

☐ was filed as PCT international application

Number \_\_\_\_\_

on \_\_\_\_\_

and was amended under PCT Article 19

on \_\_\_\_\_

I hereby state that I have reviewed and understand the contents of the above-identified specification, including the claims, as amended by any amendment referred to above.

I acknowledge the duty to disclose information which is material to the examination of this application in accordance with Title 37, Code of Federal Regulations, §1.56.

I hereby claim foreign priority benefits under Title 35, United States Code, §119 of any foreign application(s) for patent or inventor's certificate or of any PCT international application(s) designating at least one country other than the United States of America listed below and have also identified below any foreign applications(s) for patent or inventor's certificate or any PCT international application(s) designating at least one country other than the United States of America filed by me on the same subject matter having a filing date before that of the application(s) of which priority is claimed:

PRIOR FOREIGN/PCT APPLICATION(S) AND ANY PRIORITY CLAIMS UNDER 35 U.S.C. 119:

COUNTRY (if PCT, indicate "PCT")	APPLICATION NUMBER	DATE OF FILING (day, month, year)	PRIORITY CLAIMED UNDER 35 USC 119
Denmark	0128/95	February 3, 1995	<input checked="" type="checkbox"/> YES <input type="checkbox"/> NO
Denmark	1192/95	October 23, 1995	<input checked="" type="checkbox"/> YES <input type="checkbox"/> NO
Denmark	1256/95	November 10, 1995	<input checked="" type="checkbox"/> YES <input type="checkbox"/> NO
			<input type="checkbox"/> YES <input type="checkbox"/> NO
			<input type="checkbox"/> YES <input type="checkbox"/> NO
			<input type="checkbox"/> YES <input type="checkbox"/> NO

**COMBINED DECLARATION FOR PATENT APPLICATION AND POWER OF ATTORNEY**  
(Includes Reference to PCT International Applications)

Attorney's Docket Number:  
4394.400-US

I hereby claim the benefit under Title 35, United States Code §120 of any United States application(s) or PCT international application(s) designating the United States of America that is/are listed below and, insofar as the subject matter of each of the claims of this applications is not disclosed in that/those prior application(s) in the manner provided by the first paragraph of Title 35, United States Code, §112, I acknowledge the duty to disclose material information as defined in Title 37, Code of Federal Regulations, §1.56(a) which occurred between the filing date of the prior application(s) and the national or PCT international filing date of this application:

PRIOR U.S. APPLICATIONS OR PCT INTERNATIONAL APPLICATIONS DESIGNATING THE U.S. FOR BENEFIT UNDER 35 U.S.C. 120:

U.S. APPLICATIONS		STATUS (Check one)		
U.S. APPLICATION NUMBER	U.S. FILING DATE	Patented	Pending	Abandoned
08/600,908	February 13, 1996		X	
PCT APPLICATIONS DESIGNATING THE U.S.				
APPLICATION NO.	FILING DATE	US SERIAL NUMBER ASSIGNED (if any)		
PCT/DK96/00057	February 5, 1996			X




POWER OF ATTORNEY: As a named inventor, I hereby appoint the following attorney(s) and/or agent(s) to prosecute this application and transact all business in the Patent and Trademark Office connected therewith.  
Steve T. Zelson Elias J. Lambiris Cheryl H. Agris Karen A. Lowney James J. Harrington  
Reg. No. 30,335 Reg. No. 33,728 Reg. No. 34,086 Reg. No. 31,274 Reg. No. 38,711

Send Correspondence to: Steve T. Zelson, Esq.  
Novo Nordisk of North America, Inc.  
405 Lexington Avenue, Suite 6400  
New York, New York 10174-6401

Direct Telephone Calls To:  
Steve T. Zelson  
(212) 867-0123

	Full Name of Inventor	Family Name	First Given Name	Second Given Name
		Svendsen	Allan	
	Residence & Citizenship	City	State or Foreign Country	Country of Citizenship
		DK-3460 Birkerød	Denmark	Denmark
	Post Office Address	Post Office Address	City	State & Zip Code/Country
		Bakkeleddet 28	DK-3460 Birkerød	Denmark
	Full Name of Inventor	Family Name	First Given Name	Second Given Name
		Bisgård-Frantzen	Henrik	
	Residence & Citizenship	City	State or Foreign Country	Country of Citizenship
		DK-2800 Lyngby	Denmark	Denmark
	Post Office Address	Post Office Address	City	State & Zip Code/Country
		Sandkrogen 27	DK-2800 Lyngby	Denmark
3	Full Name of Inventor	Family Name	First Given Name	Second Given Name
		Borchert	Torben	Vedel
	Residence & Citizenship	City	State or Foreign Country	Country of Citizenship
		DK-2200 Copenhagen N	Denmark	Denmark
	Post Office Address	Post Office Address	City	State & Zip Code/Country
		Dagmarsgade 34, 1ejl. 17	DK-2200 Copenhagen N	Denmark

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under section 1001 of Title 18 of the United States Code, and that such willful false statements may jeopardize the validity of the application or any patent issuing thereon.

Signature of Inventor 1	Signature of Inventor 2	Signature of Inventor 3
		
Date	Date	Date
31/7 1996	5/8-96	6-8-96

CERTIFICATE OF MAILING

I hereby certify that this paper and every paper referred to therein as being enclosed is being deposited with the U.S. Postal Service as first class mail, postage prepaid, in an envelope addressed to: Commissioner of Patents & Trademarks, Washington, DC 20231.

on \_\_\_\_\_ (Date of Deposit)

Date \_\_\_\_\_ Name \_\_\_\_\_

EXPRESS MAIL CERTIFICATE

Date 6/8/99 Label No. 595859379

I hereby certify that, on the date indicated above I deposited this paper or fee with the U.S. Postal Service & that it was addressed for delivery to the Commissioner of Patents & Trademarks, Washington D.C. 20231 by "Express Mail Post Office to Addressee" service.

D B Peck  
Name (Print)

[Signature]  
Signature

Docket No.: 0776/1F216-US2

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re Application of:

Allan SVENDSEN, et al.

Serial No.: To be Assigned  
(Continuation of 08/683,838)

Art Unit:

Filed: Concurrently

Examiner:

For:  $\alpha$ -AMYLASE MUTANTS

-----  
ASSOCIATE POWER OF ATTORNEY

June 8, 1999

The Honorable Assistant Commissioner of  
Patents and Trademarks  
Washington, D.C. 20231

Sir:

The undersigned attorney of record in the above-identified application hereby appoints:

Michael J. Sweedler (No. 19,937)  
S. Peter Ludwig (No. 25,351)  
Paul Fields (No. 20,298)  
Joseph B. Lerch (No. 26,936)



Melvin C. Garner (No. 26,272)  
Ethan Horwitz (No. 27,646)  
Adda C. Gogoris (No. 29,714)  
Martin E. Goldstein (No. 20,869)  
Marc S. Gross (No. 19,604)  
Bert J. Lewen (No. 19,407)  
Henry Sternberg (No. 22,408)  
Peter C. Schechter (No. 31,662)  
Robert Schaffer (No. 31,194)  
Robert C. Sullivan, Jr. (No. 30,499)  
Ira J. Levy (No. 35,587)  
Joseph R. Robinson (No. 33,448)

located at Darby & Darby, P.C., 805 Third Avenue, New York, NY 10022-7513, as  
associate attorneys with full power to prosecute said application, and to transact all  
business in the United States Patent and Trademark Office in connection therewith.

Please forward all communications to:

Adda C. Gogoris  
Darby & Darby, P.C.  
805 Third Avenue  
New York, New York 10022-7513  
Telephone No. (212) 527-7700  
Facsimile No. (212) 753-6237

Dated: June 8, 1999

  
Elias J. Lambiris (Reg. No. 33,728)